



Complex Molecules in Space Present status and prospects with ALMA

8th-11th May 2006

Fuglsøcentret, Denmark



Including a one day meeting of the EU Network
"The Molecular Universe"



Sponsored by the EU, the Danish National Research Council, the Instrument Centre for Danish Astrophysics and the Department of Physics and Astronomy, University of Aarhus.



Table of Contents

Programme	
Monday	3
Tuesday	4
Wednesday	6
“The Molecular Universe” programme	8
Abstracts for talks arranged chronologically	
Monday	10
Tuesday	16
Wednesday	28
Abstracts for “The Molecular Universe” talks arranged chronologically	37
List of posters	50
Poster abstracts arranged in alphabetical order of first author	52
List of Participants	71
Index	76

Complex Molecules in Space

Present status and prospects with ALMA

8th – 11th May 2006, Fuglsøcentret, Denmark

Programme

Monday 8th May

12:30	Lunch	
14:30-14:45	Welcome	David Field, (University of Aarhus)
	<i>Session 1 Chair: David Field</i>	
14:45-15:20	ALMA and chemistry surveys	Ewine van Dishoeck (Leiden Observatory)
15:20-15:55	The impact of interferometry on the detection of large molecules	Tom Wilson (European Southern Observatory)
15:55-16:25	Tea/Coffee	
	<i>Session 2 Chair: John Pearson</i>	
16:25-17:00	Line surveys in Orion/SgrB2	Claudia Comito (Max-Planck-Institut für Radioastronomie, Bonn)
17:00-17:35	Molecules in Orion	Peter Schilke (Max-Planck-Institut für Radioastronomie, Bonn)
17:35-18:10	The Spectral Legacy Survey: a spectral imaging survey with JCMT	Gary Fuller (University of Manchester)
18:10-18:35	An Orion KL Survey with the Odin sub mm satellite	Henrik Olofsson (Onsala Space Observatory)
19:15	Dinner	
20:30	Poster Session	

Tuesday 9th May

8:00-9:00 Breakfast

Session 3 Chair: Eric Herbst

9:00-9:35 Exterminating the weeds in the astronomical garden Frank de Lucia (Ohio State University)

9:35-10:10 Tools for ALMA: data, databases, analysis software and the Virtual Observatory Marie-Lise Dubernet and Adam Walters (Observatory of Paris-Meudon and CNRS/Paul Sabatier Toulouse)

10:10-10:40 Coffee

Session 4 Chair: Satoshi Yamamoto

10:40-11:15 Radiative transfer tools and molecular data bases for ALMA Frederik Schöier (Stockholm Observatory)

11:15-11:50 Observations of Circumstellar Disks using the Smithsonian Millimeter Array Nagayoshi Ohashi (Academia Sinica Taipei)

11:50-12:25 Molecules in protostellar outflows: prospects for ALMA Rafael Bachiller (Observatorio Astronomico Nacional, Madrid)

12:30 Lunch

Session 5 Chair: Karl Menten

13:50-14:25 Complex organic molecules in the hot corinos of Sun-like protostars Cecilia Ceccarelli (Observatoire de Grenoble)

14:25-15:00 How are complex molecules in hot cores and corinos produced? Eric Herbst (Ohio State University)

15:00-15:25 Testing grain-surface astrochemistry through observations Suzanne Bisschop (Leiden Observatory)

15:25-16:00 Tea

Tuesday 9th May – continued

Session 6 Chair: Tom Wilson

16:00-16:35	Probing the inner 200 AU of low-mass protostars	Jes Jørgensen (Harvard-Smithsonian Centre for Astrophysics)
16:35-17:00	SMA Observations of the Class 0 Protostar Barnard 1-c	Brenda Matthews (Herzberg Institute of Astrophysics)
17:00-17:35	Molecules in massive star formation observed with high angular resolution	Henrik Beuther (Max-Planck-Institut für Astronomie, Heidelberg)
17:35-18:10	Probing high-mass star formation through different molecules	Riccardo Cesaroni (Observatorio Astrofisico di Arcetri)
19:00	Dinner	
20:30	Round table discussion: ALMA Regional Centres: how best to aid users of ALMA in preparation of applications to data analysis. Contributions from Europe, the US and Japan	Led by Tom Wilson (ESO) with contributions from Al Wootten, Satoshi Yamamoto and others.

Wednesday 10th May

08:00-9:00 Breakfast

Session 7 Chair: Ewine van Dishoeck

9:00-9:35	Chemical complexity with ALMA: looking for new molecular species near the line confusion limit	Jose Cernicharo (Instituto de Estructura de la Materia, Madrid)
-----------	--	--

9:35-10:10	From molecules to dust	Anja Andersen (University of Copenhagen)
------------	------------------------	---

10:10-10:40 Coffee

Session 8 Chair: Peter Schilke

10:40-11:15	The Cologne database for molecular spectroscopy and requirements of laboratory spectroscopy for ALMA	Holger Müller (Max-Planck-Institut für Radioastronomie, Bonn)
-------------	--	--

11:15-11:50	The Cologne THz Spectrometers: Recent Advances	Stephan Schlemmer (University of Cologne)
-------------	--	--

11:50-12:25	Molecular spectroscopy and the search for large molecules with ALMA	Sven Thorwirth (Max-Planck-Institut für Radioastronomie, Bonn)
-------------	---	---

12:40 Lunch

Session 9 Chair: Marie-Lise Dubernet

14:00-14:25	Rotational Spectra of Nucleobases	Sandra Bruenken (Harvard University)
-------------	-----------------------------------	---

14:25-14:50	Exploring an early stage of protostellar Evolution with complex organic molecules	Nami Sakai (University of Tokyo)
-------------	---	-------------------------------------

14:50-15:25	ALMA's Window on Molecular Emission: Small Ions to Complex Prebiotics	Al Wootten (National Radio Astronomy Observatory)
-------------	---	--

15:25-16:00 Tea

Wednesday 10th May – continued

Session 10 Chair: Anja Andersen

16:00-16:35	Bridging the molecular divide between molecular clouds and biology, a spectroscopist's view of detectability	John C. Pearson (Jet Propulsion Laboratory)
16:35-17:15	Summarizing the meeting	Karl Menten (Max-Planck-Institut für Radioastronomie, Bonn)
18:30	Conference Dinner	
20:30	Round table discussion: the definitive detection of large molecules.	Led by Eric Herbst with informal contributions from Al Wootten, Peter Schilke, Pepe Cernicharo, John Pearson, Sven Thorwirth and others.

Thursday 11th May

07:30-08:30 Breakfast

The Molecular Universe

Physics and chemistry of complex molecules: the laboratory approach

Interstellar PAHs

08:30-09:00	Spectroscopy of PAHs at IR/submm wavelengths	M. Vervloet (LPPM CNRS, Orsay)
09:00-09:30	High-resolution spectroscopy of large molecules	K. Demyk (PhLAM, Lille University)
09:30-10:00	Recent results on recombination of PAH cations with electrons	L. Biennier (PALMS, Rennes University)
10:00-10:30	Chemistry of interstellar PAH candidates: from space to the laboratory	C. Joblin (CESR, Toulouse University)
10:30-11:00	Coffee	

Interstellar Carbon Chains

11:00-11:30	Carbon chain spectroscopy	T. Giesen (Cologne University)
11:30-12:00	Molecular beam studies on reactions of C atoms with unsaturated hydrocarbons: a possible synthetic route to interstellar carbon chains	P. Casavecchia (Perugia University)
12:00-12:30	Intra- and intermolecular dynamics of cyanoacetylene and its complexes with helium	R. Moszynski (Varsow University)
12:30-13:00	Polyynes and cyanopolyynes : from astrochemistry to their easy synthesis in a carbon arc	F. Cataldo (Rome University)
13:00-14:00	Lunch	

Thursday 11th May - continued

Interstellar Grains: Composition and Role in Chemistry

14:00-14:30	Formation of interstellar methanol and other oxygen containing compounds: non-feasibility of gas-phase mechanisms	W. Geppert (Stockholm University)
14:30-15:00	Experimental studies of H ₂ formation on grains: overview and recent results	F. Dulieu (LERMA, University of Cergy-Pontoise)
15:00-15:30	Irradiation and thermal processing of silicates in the laboratory: a new insight in interstellar grain chemical composition	Z. Djouadi (IAS, University of Paris Sud)
15:30-15:45	Coffee	

Prebiotic/Biological Molecules

15:45-16:15	Photo- and radioracemization of chiral molecules and the radiation-induced amplification of chirality	F. Cataldo (Rome University)
16:15-16:45	Spectroscopy of complex and organic chemistry in Titan.	A. Jolly (LISA, University of Paris 12)

ALMA and chemistry surveys

Ewine van Dishoeck

Leiden University, P. O. Box 9513, Leiden, 2300 RA, The Netherlands

ALMA will revolutionize the field of astrochemistry due to its exceptional sensitivity and angular resolution down to 10 milliarcsec scales (1 AU in the nearest star- and planet-forming regions). In this talk, the prospects for ALMA will be placed in the context of recent unbiased and selected line surveys of a variety of astronomical sources, including low- and high-mass protostars, outflows, disks and envelopes around evolved stars. Special attention will be paid to those scientific questions which ALMA will be particularly well placed to address, including the origin of complex organic molecules. The need for powerful analysis tools and molecular databases to address the huge ALMA data sets will be emphasized.

The Impact of Interferometry on the Detection of Large Molecules

T. L. Wilson¹, L. E. Snyder²

¹ ALMA Division, European Southern Observatory, K-Schwarzschild-Str. 2, 85748
Garching, Germany

² Astronomy Dept., Univ. of Illinois, 1002 West Green Street, Urbana, IL 61801, USA

Up to the present, measurements with interferometers have not led to the detection of large molecules (with the notable exception of acetic acid, which was detected with the BIMA and OVRO arrays). These instruments have however contributed to our understanding of the physical conditions in which large molecules are found. The reasons for this state of affairs are investigated. We discuss new and future instruments such as the CARMA array and ALMA. We estimate how these would have contributed to the discoveries of large molecules made in the last few years.

Line Surveys in Orion and SgrB2

Claudia Comito¹, Arnaud Belloche¹, Carolin Hieret¹, Karl M. Menten¹,
Holger S.P. Müller^{1,2}, Peter Schilke¹

¹ Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, Bonn, 53121, Germany

² I. Physikalisches Institut, Universität zu Köln, Zùlpicher Str. 77, Köln, 50937, Germany

I will present an overview of the current knowledge arising from millimeter and submillimeter line surveys of the two best studied regions of massive star formation, Orion and SgrB2. Issues such as the analysis of datasets containing thousands of lines, and the detection and identification of previously undetected, complex species will also be addressed. Finally, I will provide a preview on the HEXOS Key Project for Herschel/HIFI.

Molecules in Orion

Peter Schilke

Max Planck Institute for Radioastronomie, Auf dem Högel 69, Bonn, 53121, Germany

The Spectral Legacy Survey: a spectral imaging survey with JCMT

G. Fuller¹, F. Helmich², R. Plume³ and the SLS collaboration

¹ School of Physics & Astronomy, University of Manchester, UK

² SRON, Groningen, Netherlands

³ Department of Physics & Astronomy, University of Calgary, Canada

Molecules provide the only probes which can reveal the dynamics, physics, chemistry and evolution of the cold dense regions of molecular clouds in which stars form. The Spectral Legacy Survey (SLS) is being undertaken to study the molecular inventory of a sample of molecular cloud sources. The observations will be used to investigate how the molecular properties of a source are related to the physical conditions in the gas and the source's evolutionary status.

Using the new 16 pixel 345 GHz array receiver HARP-B on the James Clerk Maxwell Telescope (JCMT), the SLS will produce spectral scans of five sources over the frequency range 330 GHz to 360 GHz. For every spectral channel throughout this frequency range of the survey will produce a fully sampled image. The five target sources to be observed have been selected to span a range of different evolutionary stages and different physical environments in molecular clouds. In this presentation I will discuss the strategy and goals of the SLS.

An Orion KL line survey with the Odin submm satellite

A.O.H. Olofsson¹, C. Persson¹, Å. Hjalmarson¹, P. Bergman, P. Bernath, J. Black, U. Frisk, W. Geppert, T. Hasegawa, N. Koning, B. Larsson, A. Lecacheux, S. Kwok, A. Nummelin, M. Olberg, Aa. Sandqvist, K. Volk and E.S. Wirström

¹ Onsala Space Observatory, SE-439 92 Onsala

We have performed a satellite based 40 GHz submm spectral survey of Orion KL in wavelength regions inaccessible from the ground due to atmospheric opacity. The frequency ranges covered are 486.4-492.1 and 541.4-577.6 GHz, and the spectral resolution is 1 MHz.

Examples of lines/molecules in the survey are 4 water isotopologues (ortho- and para-H₂¹⁶O, H₂¹⁷O, H₂¹⁸O, HDO), the J=5-4 lines from 4 carbon monoxide isotopologues (¹²CO, ¹³CO, C¹⁸O, C¹⁷O), ammonia (¹⁴NH₃ and ¹⁵NH₃), and possible designations of line features from ND, SO⁺, SH. The latter ones are not to be considered identifications since the features are very weak and it would require a number of coincidences in order to claim detections.

In total, we have identified about 300 lines from 38 molecules. An additional ~60 line features remain unidentified.

Our result also indicate that line crowding may become a problem in similar observations using future instruments such as Herschel and ALMA with higher beam filling factors and sensitivities. In particular, high-velocity line wings in the strong emission lines - often present in observations towards high-mass star forming cores - will have significant contributions from other species with high line densities such as SO₂. In order to correctly reconstruct such line wings, the strength and shape of the 'polluting' line must be accurately predicted and subtracted.

Odin is an astronomy/aeronomy satellite launched in February 2001 as a result of a collaboration between the space agencies in Sweden, France, Canada and Finland. It remains in good health and will continue operation for at least another year.

References to this work:

Olofsson, A.O.H., Persson, C., et al. 2006,
"A spectral survey of Orion KL from 487-492 and 542-577 GHz with the Odin satellite. I. The observational data." To be submitted to A&A

Persson, C., Olofsson, A.O.H. et al. 2006,
"A spectral survey of Orion KL from 487-492 and 542-577 GHz with the Odin satellite. II. Analysis and interpretation." To be submitted to A&A

Exterminating the weeds in the astronomical garden

Frank de Lucia

Dept. of Physics, The Ohio State University, Physics Research Building,
191 West Woodruff Avenue, Columbus, OH 43210, USA

The existence of unidentified features in interstellar spectra has persisted almost from the beginning of the field. In recent years the number of such lines has rapidly increased in parallel with the sensitivity and frequency range of new observational systems. Estimates of the numbers vary with source, but are typically tens of percent of all of the observed lines. Initially, they often were from 'exotic' species, such as HCO^+ , but now the origins of these lines are overwhelmingly from large molecules with dense spectra. It is expected that most of these 'weeds' are associated with previously identified interstellar species. The origin of the identification problem lies in the nature of the spectroscopic approach that has typically been used in the millimeter and submillimeter spectral region: the bootstrap observation, assignment, and theoretical prediction cycle. While this cycle is dear to the hearts of many of us, these 'weeds' arise from complex spectra involving many low lying and often interacting vibrational states. The resulting spectroscopic complexity makes the calculation of 'complete' astronomical spectra a very long-term project. In fact, the observational efforts are accelerating away from their supporting data base at a rapid pace. In this talk we will describe a purely experimental approach to this problem which does not require spectra assignment but rather relies on the observation of 'complete' spectra over a range of temperatures. This approach is enabled by the FASSST spectroscopic system and the use of Collisional Cooling cells to provide reference spectra at low temperature. Experimental and theoretical results will be presented.

Tools for ALMA: data, databases, analysis software and the Virtual Observatory

M.L. Dubernet¹ and A. Walters²

¹LERMA, Paris Observatory, 5 Place Janssen, 92195 Meudon cedex, France

²CESR, 9 Av. Du Colonel Roche, 31029 Toulouse cedex, France

Observatories such as ALMA or Herschel will provide much larger quantities of data than current instruments. Extraction of the information useful for scientific interpretation will require the use of complex numerical models including the dynamics, radiative transfer and chemistry of sources. While much effort is invested in the development of data reduction software (calibration), the tools available for high level scientific analysis of observations are poorly adapted to these new instruments and significant progress is required in this field. This is essential to maximize the scientific return from the investments made in instrumentation.

We present here the status of ongoing work on the provision of atomic and molecular reference data and on the development of automated analysis software systems.

Virtual Observatories provide a framework for gathering and providing standardized access to spectroscopic and collisional data. Recently, a data model for accessing atomic and molecular linelists has been developed and proposed as a VO standard. Another data model providing access to other types of atomic and molecular data is currently under way.

In parallel, work on databases is progressing. The current BASECOL database (<http://www.obspm.fr/basecol>) provides classical web interfaces and standardized automatic access to collisional rate coefficients for the ro-vibrational excitation of molecules by H, He, H₂ and customized spectroscopic data from JPL/CDMS. The data sets are fully documented and referenced. Visualisation tools are also provided.

CASSIS (<http://pc-126.cesr.fr/> Centre d'Analyse Scientifique de Spectres Infrarouges et Sub-millimétriques) provides interactive software allowing the user to visualize and analyse spectra taken with radiotelescopes. It accepts data files generated by CLASS software (post summer 2005) or ASCII tables (CASSIS Format), is coupled with databases and basic models, and is particularly useful for identifying lines and deducing physical parameters. The CASSIS interactive spectrum analyser is available in two modes: a dual (server + applet) standalone Java application and a web Java applet interface.

MAGIX is a collaborative project headed by the MPIfR in collaboration with several teams. This software will be a generic interface to numerical codes and will allow the user to easily fit the output of numerical models (galaxy, protostellar object, etc...) to observed data (spectrum, image, datacube, ...) in order to extract physical information. Program execution will be controlled through a user friendly graphical user interface and will not require any specific programming skills. The interface will provide the user with all the information required to understand the physics of the numerical codes and will allow the user to focus exclusively on the adjustment of the model parameters and the scientific interpretation of the data. A variety of analysis tools will also be provided to allow a critical interpretation of the modelling. As such, this interface will offer a new way to analyse data adapted to all wavelengths.

Radiative transfer tools and molecular databases for ALMA

Fredrik Schöier¹

¹ Stockholm Observatory, AlbaNova University Center, SE-10691 Stockholm, Sweden

In order to fully utilize future (sub)millimetre and infrared telescopes, such as ALMA, it is important that detailed tools are available to the general astronomical community that can match the quality of the data. Presented here is a summary of current publically available radiative transfer codes and molecular data bases. A new project named *Exciter*, aimed at providing the user with a set of advanced tools for performing both line and continuum radiative transfer calculations, will also be presented.

Observations of Circumstellar Disks using the Submillimeter Array

Nagayoshi Ohashi¹

¹ Academia Sinica Institute of Astronomy & Astrophysics, P.O. Box 23-141, Taipei 10617, Taiwan, R.O.C.

Circumstellar disks around T Tauri and Herbig Ae stars are considered to be most probable sites for planet formation, and detailed studies on them at higher angular resolutions are crucial for us to understand the mechanism of planet formation. The Submillimeter Array (SMA), recently constructed at the top of Mauna Kea, Hawaii, as a collaborative project of the Smithsonian Astronomical Observatory (SAO) and the Academia Sinica Institute of Astronomy & Astrophysics (ASIAA), is a powerful tool to observe circumstellar disks. We have been observing circumstellar disks at 230 GHz, 345 GHz, and 690 GHz bands using the SMA, revealing new aspects of circumstellar disks. In my talk, I will review observations of circumstellar disks conducted using the SMA.

Molecules in protostellar outflows: prospects with ALMA

Rafael Bachiller¹

¹ Observatorio Astronómico Nacional, IGN, c/ Alfonso XII, 3, 28014 Madrid, Spain

The propagation of a supersonic protostellar outflow through its surrounding medium happens via shock waves. The rapid heating and compression of the gas trigger different microscopic processes – such as molecular dissociation, endothermic reactions, ice sublimation, and dust grain disruption –, which do not operate in the unperturbed gas, generating peculiar chemical anomalies.

We review, from an observational point of view, the influence of shocks on chemistry emphasizing recent observations of outflows from Class 0 sources with favorable orientation in the sky (i.e. high inclination with respect to the line of sight). With less confusion than that found around massive outflows, the shocked regions of low-mass high-collimation outflows, often adopt the form of well-defined bows which are well separated spatially with respect to the quiescent gas. The behaviour of key molecules such as SiO, CH₃OH, H₂CO, HCO⁺, CN, etc, is summarized. Particular attention is paid to the chemistry of sulfur bearing molecules (H₂S, OCS, CS, SO, SO₂) since it has been recently proposed to be a potential tool to construct chemical clocks to date outflows (and hence their protostellar driving sources). We also present recent results of a comprehensive chemical survey aimed to study the evolution of young protostellar outflows (Ph. D. work of J. Santiago). It appears that the peculiarities of the chemical behaviour together with other structural parameters such as the presence of extremely high velocity features (“molecular bullets”), the collimation factors, and the mechanical power efficiencies, can be used to produce a rough classification of the observed outflows and to suggest a time evolutionary sequence.

We finally provide some prospects with the future ALMA observations. The superb sub-arcsecond angular resolution together with the high sensitivity in a variety of molecular lines will be particularly useful to study the outflow launching regions, the details of the entrainment mechanisms, and the abundance of very rare molecules which are expected to be created by the shocks.

The complex organic molecules in the hot corinos of Sun-like protostars

Cecilia Ceccarelli

Laboratoire d'AstrOphysique de Grenoble, 414 rue de la Piscine BP53, Grenoble, 38041,
France

Sun-like stars form from cold and dense condensations inside molecular clouds. During the first phases, heavy element bearing molecules freeze-out onto the dust grains forming mantle of water ices with traces of other molecules. When the collapse finally starts, the object at the center of the condensation increases in mass and in luminosity. The central regions are consequently heated up, and can reach the temperature at which the icy grain mantles sublime, injecting back the molecules accreted during the pre-collapse phase, along with those created on the grain surfaces. These central regions have been observed in a few low mass protostars, and have been named “hot corinos” to distinguish them from the massive cousins “hot cores”. The radii of the hot corinos are less than about 150 AU and the densities are larger than about 10^7 cm^{-3} . Although very few cases have been studied in detail, there is a strong evidence that hot corinos are a common phase in the life of a Sun-like protostar. The observations also suggest that complex organic molecules are very abundant in hot corinos, equating the abundance of formaldehyde and/or methanol. Remarkably, complex organic molecules seem to be more abundant in hot corinos than in massive hot cores, when compared to methanol and formaldehyde in both classes of objects. It is unclear yet whether this is due to a different temperature in the hot cores and corinos, or rather totally to a substantial difference in the composition of the grain mantles. Based on the observations so far available, it seems likely that the observed complex organic molecules are mostly formed on the grain surfaces. In this contribution I will review this topics and discuss how ALMA will represent a revolutionary instrument for these studies, allowing to detect and image the hot corinos, whose predicted typical sizes are out of reach of the current interferometers.

How Are Complex Molecules Produced In Hot Cores And Corinos?

Eric Herbst

Departments of Physics, Astronomy, and Chemistry, The Ohio State University,
Columbus, OH 43210 USA

Hot cores and corinos are warm (100-300 K) regions in the vicinity of protostars where a large variety of hydrogen-rich organic molecules are found. Until recently, it had been thought that the molecules were produced in the gas mainly from the precursor molecule methanol, which had been synthesized on the surfaces of dust particles in a previous colder eon before the onset of star formation. Once evaporated into the gas as temperatures increase, the methanol is protonated, leading to a rich ion-molecule chemistry. Unfortunately, some of the reactions thought to form more complex molecules have now been measured not to occur. We suggest another method in which larger molecules can be produced – in this scenario the molecules are produced as the temperatures increase during star formation and not only afterwards. The basic idea is that a distinct chemistry can occur on the surfaces of dust particles warmed to higher temperatures than 20 K, where H atoms are not as important in the chemistry as at lower temperatures and heavy radicals are able to diffuse and react with one another. As the surface temperatures increase to 100 K, the newly formed species are evaporated and change the nature of the gas-phase chemistry.. The specific case of methyl formate, a common and abundant species in hot cores, is looked at in some detail.

Testing grain-surface chemistry in massive hot-core regions

Suzanne E. Bisschop¹, Jes K. Jørgensen², Ewine F. van Dishoeck¹

¹ Leiden Observatory, Postbus 9513, 2300 RA Leiden, Netherlands

² Harvard-Smithsonian Center for Astrophysics, 60 Garden Street MS42, Cambridge, MA 02138, USA

Many complex organic molecules, such as CH₃OH, CH₃OCH₃ and CH₃CH₂CN, have been detected in star-forming regions. Studying their chemistry is important, since these species may eventually be incorporated into circum-stellar disks and thus become part of the material from which future planetary systems are made. Two scenarios for their formation have been proposed: grain-surface chemistry and high-temperature gas-phase reactions (see e.g., van Dishoeck & Hogerheijde, 1999 for a review). In the former case, grain-surface reactions during the cold pre-stellar and proto-stellar phases are thought to lead to the formation of hydrogenated molecules, which subsequently evaporate into the gas when the young star heats its surroundings. In the latter case, high-temperature gas-phase reactions involving evaporated molecules produce complex organic species like CH₃OCH₃ (Charnley et al., 1992). The resulting molecules from these two schemes are also known as "first generation" and "second generation" species. Currently, it is very difficult to distinguish between these two scenarios since the rates of even some of the most basic surface reactions are not known.

Deep searches for a set of molecules, that are thought to have formed on grain surfaces, have been performed for 7 high mass Young Stellar Objects with the James Clerk Maxwell Telescope (JCMT). We detect emission from H₂CO, CH₃OH, CH₂CO, CH₃CHO, C₂H₅OH, HCOOH, HNCO and NH₂CHO. Additionally, molecules that are thought to form from gas phase reactions with CH₃OH, such as CH₃OCH₃, and CH₃OCHO, have been detected. CH₃CHO is not detected with the JCMT for any source; it was previously detected by Ikeda et al. (2002) with rotational temperatures of only 20-30 K, which is much lower than the lowest rotational transition that can be probed with the JCMT at 71 K. This is also confirmed by our detections of CH₃CHO transitions with energy levels below 45 K in N7538 IRS1, and W3(H₂O) with IRAM, two of the sources from our JCMT sample. The resulting rotational temperatures, derived from rotation diagrams, are about 100 K or higher for most target molecules, but are significantly lower for CH₂CO, and CH₃CHO, implying that these are not present in the same region.

The derived column densities vary by more than an order of magnitude between the different sources. The relative column densities are however very similar for some molecular species, e.g., H₂CO, CH₃OH, C₂H₅OH, CH₃OCHO, CH₃OCH₃. Surprisingly, this includes both molecules that are thought to form due to grain-surface reactions and gas phase chemistry. The column densities of the nitrogen-bearing species HNCO and NH₂CHO behave very similar as well, but peak for other sources compared to the oxygen-bearing species. Abundances are derived by comparing to H₂ column densities estimated from dust radiative transfer models. Strong correlations are found between those species that show very similar column density variations suggesting that they are chemically related through their formation mechanisms.

Probing the Inner 200 AU of Low-Mass Protostars with the Submillimeter Array and Spitzer Space Telescope

Jes K. Jørgensen¹, and the PROSAC² and c2d³ teams

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, MS42,
Cambridge, MA 02138, USA

² Protostellar Submillimeter Array Campaign

³ “Cores to Disks” Spitzer Space Telescope Legacy program

One of the more exciting recent developments in astrochemistry is the detection of complex organic molecules in the envelopes around low-mass protostars. Studies of this chemistry are important because part of this material may eventually be incorporated in circumstellar disks and new planetary systems. Low-mass protostars are now known to contain “hot corinos” similar to the hot cores observed for high-mass protostars, i.e., warm ($T \sim 100$ K), dense ($n_{\text{H}_2} \sim 10^7 \text{ cm}^{-3}$) regions where icy grain mantles evaporate, injecting organic molecules into the gas-phase where they can further react to form more complex species (e.g., Ceccarelli et al. 2000, Schöier et al. 2002, Cazaux et al. 2003).

The origin of this warm gas with complex molecules is strongly debated, however. In addition to evaporation of ices following passive heating of the inner envelope by the protostar, also interactions of the outflow with the inner envelope can be important. For example, H_2CO and CH_3OH - often taken as signposts for hot corinos - have been shown to be affected by shocks in a number of protostellar outflows on both large scales and in the immediate vicinity of low-mass protostars (e.g., Bachiller et al. 1995, Jørgensen et al. 2005). Also, estimates of molecular abundance variations rely on a good understanding on the envelope physical structure, in particular its density and temperature, down to smallest scales. Estimates of these parameters often rely on extrapolations of the structure from the outermost regions and may break down on small scales where also the protoplanetary disks are formed.

In this talk I will discuss some of the observational constraints on the structure of the inner envelope and the chemistry in the immediate environments of low-mass protostars. In particular, I will focus on a survey of a large sample of low-mass protostars using the Submillimeter Array (SMA). Those observations directly image high excitation transitions probing the warm and dense gas in the protostellar envelopes and make it possible to trace the physical and chemical structure down to scales of a few hundred AU. In particular, the importance of shocks on those scales is apparent in the data. The SMA observations also place strong constraints on the existence and sizes of circumstellar disks on small scales.

I will also discuss recent Spitzer Space Telescope observations from the c2d legacy team that provide further clues to the structure of the innermost regions of protostellar envelopes. In particular, the mid-IR detections of the heavily obscured protostars suggest the existence of inner cavities within which material is falling toward the central circumstellar disks. The Spitzer observations also provide direct images of the warm gas in the shocks through imaging H_2 rotational emission – and show clear correlations with the strong CH_3OH emission on scales from a few hundred to thousand AU scales.

New SMA and JCMT results on the remarkable class 0 source Barnard 1c

Brenda Matthews¹, Antonio Crapsi², Michiel Hogerheijde²,
Jes Jørgensen³, Edwin Bergin⁴

¹ Herzberg Institute of Astrophysics, National Research Council of Canada,
5071 West Saanich Road, Victoria, BC, V9A 3T3, Canada

² Leiden Observatory, P.O. Box 9513, 2300 RA, Leiden, Netherlands

³ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street MS42,
Cambridge, MA, 03138

⁴ Department of Astronomy, University of Michigan, 825 Dennison Building,
500 Church Street, Ann Arbor, MI, 48109

We report recent results from molecular line studies on the Smithsonian Submillimeter Array and the James Clerk Maxwell Telescope toward the recently identified class 0 object, Barnard 1c. This source, first identified in polarimetry mapping with SCUBA (Matthews & Wilson 2002), was primarily of interest due to its unique polarization properties compared to all other low mass cores. In B1c, there is evidence that grains are well aligned by the magnetic field all the way to the core centre. Followup studies of the core interior were undertaken to understand why the polarization properties of this core are so unique. B1c is now known to drive a spectacular precessing outflow as imaged by Spitzer (Jørgensen et al. 2006) associated with a CO outflow as detected with the SMA (see Figure 1) and exhibits a rotating molecular torus on the order of 1000 AU encompassing a central heated cavity as detected with the BIMA array (Figure 1, Matthews et al. 2006). We will present our most recent results from the SMA and JCMT which reveal the flattening of the central heated core and the evidence for strong deuteration in this object, which can be linked to strong coupling of the gas to the magnetic field.




Figure 1: *Left*: The CO outflow is strongly detected in CO 2-1 emission from the SMA. The greyscale shows the 1.3 mm emission. As detected in the CO 1-0 emission, we see evidence of a shift from blue (dark grey contours) to red-shifted (light grey contours) emission where the Spitzer data reveal a change in the direction of the outflow. *Right*: The greyscale shows the first moment of a single hyperfine component of N_2H^+ 1-0 emission from B1c; the rotation from roughly north to south is evident. The contours show the zeroth moment; two peaks of N_2H^+ 1-0 emission are roughly symmetric about the continuum peak, indicated by the cross. The outflow axis and alignment of the peaks are indicated by the dash-dot and dashed grey lines, respectively.

Jørgensen, J., et al. 2006, accepted to the *Astrophysical Journal*

Matthews, B., Hogerheijde, M., Bergin, E., & Jørgensen, J. 2006, submitted to the *Astrophysical Journal*

Matthews, B., & Wilson, C. 2002, *ApJ*, 574, 822

Molecules in massive star formation observed with high angular resolution

Henrik Beuther

¹ Max-Planck-Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany

One of the open promises in (high-mass) star formation research is to develop chemical clocks allowing to better determine ages and evolutionary sequences. While some theoretical models exist, the observational database to calibrate the models against is still poor. This is specifically the case for high-angular-resolution observations resolving the various sub-sources within the evolving massive proto-clusters. I will present recent spectral line observations from (sub)mm wavelength Interferometers toward massive star-forming regions of various evolutionary stages, outlining the present state of research and the current observational possibilities. A short outlook on the progress expected by ALMA will be given as well.

Probing high-mass star formation through different molecules

Riccardo Cesaroni

INAF-Osservatorio Astrofisico di Arcetri, Largo Fermi 5, Firenze, I-50125, Italy

Star formation proceeds through inside-out collapse of molecular cores, with accretion onto a compact kernel, the protostar. While this certainly holds for low-mass stars, the same mechanism cannot be trivially applied to high-mass ones (i.e. stars in excess of about $8 M_{\text{sun}}$), as their powerful radiation pressure could halt and even reverse the infall. Observations can shed light on this issue by analysing the structure and kinematics of the gas from the scale of the molecular cores hosting OB star formation (0.1 pc) to the regions where accretion is expected to dominate (<1000 AU). After briefly reviewing the characteristics of OB star forming regions, I will show how different molecular transitions can be used to probe different parts of the star forming environment. In particular I will focus on the detection of rotation and infall towards high-mass young stellar objects, which will be illustrated in detail by means of detailed studies of selected objects. Application of the most powerful observational techniques available to date suggests that only the combination of sub-arcsec angular resolution, high sensitivity, and - last but not least - wide spectral coverage provided by ALMA will make it possible to establish the mechanism by which massive stars form.

**Chemical complexity with ALMA: looking for new molecular species
near the line confusion limit**

Jose Cernicharo

DAMIR, Instituto de Estructura de la Materia, C/Serrano 121, Madrid 28006, Spain

From Molecules to Dust

Anja C. Andersen¹, Susanne Höfner²

¹ Dark Cosmology Centre, Niels Bohr Institute, Juliane Maries Vej 30, DK-2100 Copenhagen, Denmark, email: anja@dark-cosmology.dk

² Department of Astronomy & Space Physics, Uppsala University, Box 515, SE-75120 Uppsala, Sweden, email: hoefner@astro.ku.dk

The dust injection from Asymptotic Giant Branch (AGB) stars into the interstellar medium very likely plays a key role for the dust formation in a galaxy. Determining the amount of dust injected by AGB stars into the interstellar medium as well as the composition of the dust is a pre-requisite for any attempt to model the evolution of interstellar matter. The dust in the interstellar medium is assumed to play a significant role for the formation of molecules, in particular for H₂, whose abundance relative to H atoms in the ISM cannot be accounted for by gas-phase reactions and methanol which involve heterogeneous reactions on grain surfaces or in the icy mantles.

In most dusty objects the grain component is intimately interwoven with the gas component and the radiation field and therefore strongly affects the thermodynamical, the hydro-dynamical and the chemical structure of the object.

Dust is formed by a series of chemical reactions in which atoms or molecules from the gas phase combine to clusters of increasing size. The molecular composition of the gas phase determines which atoms and molecules are available for the cluster formation and grain growth. This can lead to a layered structure in a circumstellar envelope. Dust formation alters the atmospheric structure which significantly influences the local and global circumstellar envelope dynamics and the detailed time-dependent spectral appearance.

In the pulsating atmospheres of AGB stars the condensation of dust grains is strongly connected with the shock wave structure of the circumstellar envelope, usually triggered by a two-step mechanism. First there is nucleation of critical clusters from molecules and this is followed by growth to macroscopic dust grains. Grain growth will usually proceed far from equilibrium and it is necessary to use a time-dependent description of grain growth to determine the degree of condensation and other relevant properties of the dust.

We are currently investigating how the predicted wind properties of AGB stars are influenced by the choice of micro-physical dust parameters and the stellar metallicity.

The Cologne database for molecular spectroscopy and requirements of laboratory spectroscopy for ALMA

Holger Müller

Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany
I. Physikalisches Institut, Universität zu Köln, Zùlpicher Str. 77, 50937 Köln, Germany

The internet browser-accessible Cologne Database for Molecular Spectroscopy, CDMS, contains as one main part a catalogue of rotational and, to a lesser extent, rovibrational transition frequencies of molecular species that have been or may be identified in the interstellar medium, circumstellar envelopes, or planetary atmospheres. The entries have been created by fitting critically evaluated experimental data, mostly from laboratory experiments, to suitable Hamiltonian models to ensure entries as reliable as possible. Particular emphasis has been put on supplying data in the (sub-) millimeter and terahertz ranges for telescopes such as APEX and the SMA as well as for the upcoming Herschel mission and ALMA. Furthermore, the CDMS provides data with increased accuracy for many species. Separate entries are provided for minor isotopic species or for molecules in excited vibrational states as far as appropriate. The database is available online free of charge through a link on the KOSMA web-site <http://www.ph1.uni-koeln.de/> or via the short-cut <http://www.cdms.de/>.

The catalogue is updated continuously. As of April 2006, the catalogue contains more than 360 species of astro-physical, astrochemical, and planetary interest. The basic features of the CDMS as well as recent and future activities in the field of complex molecules will be discussed.

A second part of the talk will be devoted to some issues concerning the detectability of complex molecules in the interstellar medium. In this context, a few selected examples of often not so complex molecules will be given that need further laboratory spectroscopic investigations.

The Cologne THz Spectrometers: Recent Advances and Perspectives

Stephan Schlemmer, Thomas Giesen and Frank Lewen

I. Physikalisches Institut, Universität zu Köln, 50937 Köln, Germany

Technical developments of recent years have lead to an extension of the accessible spectral range towards shorter wavelengths and into the so called Terahertz region. Many molecules, especially those of astrophysical importance, have characteristic spectral features at Terahertz frequencies, e. g., the pure rotational transitions of light hydrides, XH and XH₂ (X=C, O, N, S) or the low lying bending modes of carbon containing chain and ring molecules.

The Cologne group uses backward wave oscillators (BWOs) and a far-infrared laser sideband system to cover the spectral range up to 2 THz at very high resolution. New superlattice frequency multipliers have extended the accessible spectral range up to 2.7 THz. The spectrometers are combined with supersonic jet and trap apparatuses where unstable and highly reactive molecules are produced in a cold jet expansion or storage device. We report on our latest results and new technical developments on Terahertz radiation sources as well as laser induced reactions, a novel method for spectroscopy with ultimate sensitivity.

All molecular parameters and line catalogues are made available via the Cologne Database for Molecular Spectroscopy (CDMS) for future observations.

Molecular spectroscopy and the search for (large) molecules with ALMA

S. Thorwirth

Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany

The Atacama Large Millimeter Array, ALMA, will enable the investigation of astronomical molecules at high sensitivity and unique spatial resolution. It will also be a prime facility to search for new molecules in space. The analysis of these observational data – in particular from line rich sources - will be tedious and largely dependent on the availability and accuracy of corresponding laboratory data. Selected examples of plausible candidates for astronomical detection will be given.

The Rotational Spectra of Nucleobases

Sandra Brünken¹, Patrick Thaddeus¹, Ronald D. Brown², Peter D. Godfrey²

¹ Harvard Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02155, USA

² Monash University, Wellington Road, Victoria 3800, Australia

The nucleic acids bases (nucleobases) are one of the major constituents of DNA and RNA, and they represent next to amino acids and sugars another class of prebiotic molecules of potential astrophysical interest. Nucleobases are nitrogen containing heterocyclic compounds and can be divided into two groups according to their molecular structure: uracil, cytosine and thymine consist of a single six-membered ring and belong to the pyrimidine derivatives, whereas adenine and guanine are purine derivatives, consisting of a six-membered ring fused to a five-membered ring. They exhibit a dense rotational spectrum and are fairly polar with dipole moments ranging from 2-5 Debye.

In this talk I am going to give a summary of the available experimental data on the rotational spectra of the five nucleobases [1] and present recent results on uracil, the smallest member of the class. The existing dataset on this molecule has been extended both to the astrophysical important frequency range between 70-100 GHz, where the strongest lines are expected for interstellar sources with excitation temperatures around 10 K, and towards the microwave region, which allowed for the analysis of its complex nuclear hyperfine structure.

Additionally, possible sources for its astrophysical detection will be discussed.




Figure 1: Chemical structure of the five nucleobases.

References

- [1] Brown, R. D., Godfrey, P., McNaughton, D., Pierlot, A. P., *J. Am. Chem. Soc.* 110 (1988) 2329–2330;
 Brown, R. D., Godfrey, P., McNaughton, D., Pierlot, A. P., *J. Am. Chem. Soc.* 111 (1989) 2308-2310;
 Brown, R. D., Godfrey, P., McNaughton, D., Pierlot, A. P., *Chemical Physics Letters* 156 (1989) 61-63;
 Brown, R. D., Godfrey, P., McNaughton, D., Pierlot, A. P., *J. Chem. Soc., Chem. Commun.* 1 (1989) 37-38.

Exploring an Early Stage of Protostellar Evolution with Complex Organic Molecules

Nami Sakai¹, Takeshi Sakai², Satoshi Yamamoto¹

¹nami@taurus.phys.s.u-tokyo.ac.jp, yamamoto@taurus.phys.s.u-tokyo.ac.jp,

Department of Physics, The University of Tokyo,
Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan

²sakai@nro.nao.ac.jp, Nobeyama Radio Observatory,
Minamimaki, Minamisaku, Nagano 384-1305, Japan

Complex organic molecules such as HCOOCH_3 , CH_3OCH_3 , and $\text{C}_2\text{H}_5\text{CN}$ are known to exist in hot cores of massive star forming regions, and are key molecules to diagnose in the densest part of star forming regions. However, recently these complex organic molecules have also been detected toward a few low mass star forming regions, IRAS16293-2422, NGC1333IRS4A, NGC1333IRAS2A and NGC1333IRAS4B (Cazaux. et al. 2003, Bottinelli et al. 2004, Jørgensen et al. 2005, Sakai et al. 2006). Figure 1 shows our detection of HCOOCH_3 toward NGC1333IRAS4B at 89.3 GHz with Nobeyama 45 m telescope. These results clearly established the existence of "hot cores" even in a low mass star forming region. Complex organic molecules are produced by grain mantle evaporation and subsequent gas phase reactions just after the onset of star formation. Furthermore these sources are all the class 0 protostars, indicating that the complex organic molecules can exist in the early stage of protostellar evolution. Particularly NGC1333IRAS4B is reported to be extremely young (\sim a few 100 yr) according to the dynamical age of the molecular outflow (Choi 2001). Since large organic molecules are generally absent in starless cores (Kaifu et al. 2004), their spectral lines could be used as a novel tracer for detecting an onset of star formation.

Very recently we have found a similar trend in a massive star forming region, NGC2264, located 800 pc from the Sun. Figure 2 shows integrated intensity map of CS(7-6) (Shreyer et al.1997). In this region, IRS1 is a young stellar object with the brightest IR source. But we were not able to detect the complex organic molecules there. In contrast, we have detected the lines of HCOOCH_3 in this southern region including MMS3, which is reported to be younger than IRS1 (Ward-Thompson et al. 2000).




Fig. 1 Spectral line profile of HCOOCH_3 toward NGC1333 IRAS 4B (81, 8 - 71, 7 E and A).




Fig. 2 Integrated intensity map of CS(7-6) in NGC2264 IRS1 region. (Shreyer et al.1997 JCMT)

REFERENCES

- Bottinelli, S., et al. 2004, ApJ, 615, 354
 Cazaux, S., et al. 2003, ApJ, 593, L51
 Choi, M. 2001, ApJ, 553, 219
 Jørgensen, J. K. et al. 2005, ApJ, 632, 973
 Kaifu, N., et al. 2004, PASJ, 56, 69
 Sakai, N., Sakai, T., Yamamoto, S. 2006, PASJ, 58, L15
 Schreyer, K., Helmich, F. P., van Dishoek, E. F., & Henning, Th. 1997, A&A, 326, 347
 Ward-Thompson, D., Zylka, R., Mezger, P. G., & Sievers, A. W. 2000, A&A, 355, 1122

ALMA's Window on Molecular Emission: Small Ions to Complex Prebiotics

Al Wootten

National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, Virginia 22901, USA

The Atacama Large Millimeter Array (ALMA¹) is a large international telescope project under construction in northern Chile on a site at Chajnantor of 5 km elevation. The excellent atmospheric transmission at that site in the millimeter and sub-millimeter wavelength ranges will allow ALMA to provide detailed images of the sources of the Cosmic Microwave Background and the Cosmic Far-Infrared Background radiation, near the wavelengths of the two strongest peaks in the spectral energy distribution of the Universe. ALMA's images will contain all of the flux in the imaged field through the use of two parts: (1) the "12m Array", composed of up to sixty-four 12-meter antennas that can be placed on 186 different stations for baselines up to 18 km (see Table 1) and (2) the "Atacama Compact Array", or ACA, that consists of twelve 7 meter telescopes placed in compact configurations and four 12 meter telescopes for measuring source total power. At the shortest planned wavelength, $\lambda=0.3$ mm, and longest baseline, the angular resolution will be 0.004". The receivers use superconducting (SIS) mixers which, in combination with the excellent site transparency (median $\tau_{225}=0.061$) and the large array collecting area (8152 m²), will provide a sensitivity at 1mm wavelength of 1 mJy in a few seconds for average atmospheric conditions; this is more than two orders of magnitude better than any array operating today. Complete frequency coverage from 10mm to 0.3mm will offer unprecedented access to the spectrum of interstellar molecules at this high sensitivity. At first light for the ALMA project the 6 highest priority receiver bands will be installed (see Table), each observing both polarizations with a bandwidth of 8 GHz; construction of the additional four bands will be accomplished during the Operations phase of ALMA. The ability to map the spatial distribution of large molecules at high sensitivity and resolution and in many transitions will revolutionize our understanding of the mechanisms of formation of these species in the interstellar medium.

ALMA Front End Key Specifications (and Preliminary Results)

All Bands Offer Dual Linear Polarization; 8GHz Output per Polarization

Each 12m Antenna will also be equipped with a 183 GHz Water Vapor Radiometer

ALMA BAND	FREQUENCY (GHZ)	T _{RX} 80% OF BAND	MIXING SCHEME	RECEIVER TECHNOLOGY
1	31.3-45.	17	USB	HEMT
2	67-90	30	LSB	HEMT
3	84-116	37 (35)	2SB	SIS
4	125-169	51	2SB	SIS
5 ^a	163-211	65	2SB	SIS
6	211-270	83 (40)	2SB	SIS
7	270-373*	147 (80)	2SB	SIS
8	385-500	98	DSB	SIS
9	602-720	175 (120)	DSB	SIS
10 ^b	787-950	230	DSB	SIS

*between 370 – 373 GHz Trx is less than 300 K

a Six receivers in construction phase.

b Full complement to be finished after construction phase.

¹ ALMA is an equal partnership between Europe and North America, in cooperation with the Republic of Chile, and is funded in North America by the U.S. National Science Foundation (NSF) in cooperation with the National Research Council of Canada (NRC), and in Europe by the European Southern Observatory (ESO) and Spain. In the bilateral project, ALMA construction and operations are led on behalf of North America by the National Radio Astronomy Observatory (NRAO), which is managed by Associated Universities, Inc. (AUI), and on behalf of Europe by ESO. In addition, Japan has also entered the ALMA project.

Bridging the molecular divide between molecular clouds and biology, a spectroscopist's view of detectability

John Pearson

Jet Propulsion Laboratory, California Institute of Technology, USA

More than 130 molecules have been identified in the interstellar medium and numerous unidentified features widely believed to molecules transcend the spectrum from radio through the near ultraviolet. The identified molecules include examples of all the functional groups of organic chemistry suggesting that interstellar processing is partly if not completely responsible for the chemistry leading to life. Constraining the interstellar role in the evolution of biology is a central question in understanding the likelihood of life elsewhere. ALMA, Herschel and other large telescope facilities have a scientific mandate to bridge the gap between the relatively simple molecules already identified and the macromolecules of biology. The three micron bands provide a compelling case that more complex molecules do exist in the ISM, but there is no identification of any specific carriers. The physics and chemistry of large molecules make detection and identification difficult due to large partition functions, low volatility, and large rotational constants. However, there are many large molecules that have favorable ro-vibrational characteristics for detection. Identifying the more likely candidate molecules and eliminating the contamination lines from known molecules is required for convincing detections. Both cases will require a focused investment in laboratory work that transcends telescope facilities and wavelengths.

Spectroscopy of PAHs at IR/submm wavelengths

M. Vervloet¹, O. Pirali²¹ LPPM, CNRS, Bât 350, Université Paris-Sud, Orsay, 91405, France² FOM Institute for Plasma Physics, Edisonbaan 14, Nieuwegein, 3439MN, The Netherlands

In the last few years, we have developed an experimental set-up in order to record gas phase emission and absorption spectra of small PAHs in the far-infrared region. Their spectra have been obtained by means of a high resolution Fourier transform spectrometer. Thermal emission of PAH vapour at about 450 °K was emitted from an electric furnace loaded with solid PAH sample following the method described in ref. (1) and (2). Emission vibrational bands of naphthalene, anthracene, phenanthrene, chrysene and pyrene have been recorded in the spectral range 50-700 cm⁻¹. A Tight-Binding Molecular Dynamics approach allows the prediction of their spectral positions (3). Absorption spectroscopy was carried out on naphthalene at 300 °K, in the same spectral range, at higher resolution, by using a multiple reflection cell. The resulting spectrum can be straightforwardly compared with its 300 ° K emission spectrum recorded by using an appropriate set-up. Thanks to the high resolution, some features observed in the absorption spectrum of naphthalene can be tentatively assigned.

These far-infrared measurements provide information on the entire molecular skeleton of each PAH and characterise each species. It is hoped that they will help in the identification of some spectral astrophysical structures.

1 O. Pirali, PhD thesis, Université Paris XI, Centre Scientifique d'Orsay, (2004).

2 O. Pirali, M. Vervloet, Chem. Phys. Lett. (in press).

3 O. Pirali, Nguyen-Thi Van-Oanh, P. Parneix, M. Vervloet, P. Bréchnignac (submitted).

High-resolution spectroscopy of large molecules

K. Demyk

PhLAM, Université Lille 1, Bat. P5 Lille 59655, France

Recent results on the recombination of PAH cations with electrons

Ludovic Biennier, Mohamed Alsayed-Ali, Oldrich Novotny¹, Daniel Travers, and Bertrand Rowe

Laboratoire de Physique des Atomes, Lasers, Molécules et Surfaces, UMR 6627 CNRS et Université de Rennes 1, Rennes, France

¹ Also from: Charles University, Faculty of Mathematics and Physics, Dept. of Electronics and Vacuum Physics, V Holešovičkách 2, Prague 8, Czech Republic

Mid-infrared astronomical observations suggest that Polycyclic Aromatic Hydrocarbons (PAHs) are ubiquitous in the Universe. However, various aspects of the chemistry of PAHs remain not well understood. This is partly due to the scarcity of laboratory data, especially in the gas phase, as it is challenging to investigate the spectroscopy and kinetics of free molecular species of such low vapour pressure.

As PAHs molecules have low ionization potential they can easily be ionized in space, either by charge exchange or by photoionization. The cations can then react with neutral atoms or molecules or recombine with electrons. Among all the processes that take place in the interstellar medium, recombination may play an important role in the destruction of PAH ions.

In our laboratory, we have recently developed a new method dubbed F/API (Flowing Afterglow with PhotoIons) for measuring the recombination rate of large ionized species such as PAHs with electrons in a flowing afterglow at room temperature. A very small amount of neutral PAH molecules is injected into the afterglow plasma by evaporation from a plate coated with the PAH solid sample. PAH ions are then produced by photoionization of the parent molecule using a pulsed VUV laser (157nm). The laser beam is oriented along the flow-tube and so a constant spatial concentration of photoions is obtained. The electron concentration along the flow-tube is measured by means of a movable Langmuir probe. Ion concentration decay in time is measured at a fixed position using a quadrupole mass spectrometer which is triggered by the laser pulse. The recombination rate is then obtained from the time profile of the photoions which depend on the electron number density present in excess. The F/API method has been successfully applied to the measurement of the recombination rate of anthracene, pyrene, azulene and acenaphthene cations^{1,2}. Our measurements reveal a definite trend of increasing rate with the size of the PAH (number of carbon atoms).

However, the F/API method is limited by the vapour pressure of PAHs which strongly decrease with size. To overcome this limitation, we are currently working on a new implementation to produce the PAH ions and extend our investigations to larger species. Feasibility studies suggest that large PAHs may be efficiently photo-desorbed in situ from a pellet positioned in the flow by a first laser ($\lambda=266$ nm) and then ionized by a second high energy laser ($\lambda=157$ nm). From the history of the packet of PAH ions and photoelectrons, reconstructed by time resolved Langmuir probe and mass spectrometry measurements, and guided by numerical simulations, we can access the recombination coefficient. Preliminary results suggest that the method may hold a great potential for the measurement of the recombination rate coefficient of large condensable species such as PAHs.

This work is supported by the Programme de Physique et Chimie du Milieu Interstellaire.

References:

- 1 O. Novotny, B. Sivaraman, C. Rebrion-Rowe, D. Travers, L. Biennier, J. B. A. Mitchell and B. R. Rowe, *J. Chem. Phys.*, 2005, 123, 104303.
- 2 L. Biennier, M. Alsayed-Ali, A. Foutel-Richard, O. Novotny, S. Carles, C. Rebrion-Rowe and B.R. Rowe, *Faraday Discussion* (in press)

Chemistry of interstellar PAH candidates : from space to the laboratory

C. Joblin¹, N. Bruneleau, A. Simon & M. Armengaud, P. Frabel, L. Noguès

¹ CESR, CNRS-Univ. Toulouse III &OMP, Toulouse, France

Polycyclic Aromatic Hydrocarbons (PAHs) are the best candidates to account for the Aromatic Infrared Bands (AIBs) observed in emission between 3.3 and 12.7 μm in many astronomical environments irradiated by UV photons. Despite a wealth of observational data, no individual molecule could be identified so far. The true nature of interstellar PAHs has therefore to be questioned, considering the formation mechanism of these species, the processing by UV photons and the reactivity with gas-phase species. These studies require a multidisciplinary approach combining observations, fundamental data from theory or experiment, and astrophysical models.

One of the challenges for laboratory experiments is to approach the physical conditions that prevail in interstellar space: cold environment (10-50 K), absence of collisions on long timescales and presence of stellar UV photons ($h\nu < 13.6$ eV) at the surface of clouds. PIRENEA, "*Piège à Ions pour la Recherche et l'Etude de Nouvelles Espèces Astrochimiques*", is an experimental set-up which has been developed according to these specifications. Using the trapping, and the mass spectrometry analysis of an ion cyclotron resonance (ICR) cell that is connected to cryogenic shields, novel studies can be performed on processes which occur on long timescales and which are of interstellar relevance: IR radiative cooling, photodissociation in low energy channels, radiative association.

Results will be presented on the photodissociation of PAH species and their reactivity with small molecules. These studies extend now to nano-grains such as PAH clusters and PAH-iron complexes.

Carbon chain spectroscopy

T. Giesen

Physikalisches Institut, University of Cologne, Zulpicher Str 77, Cologne 50937,
Germany

Carbon chains represent the dominant structural theme of the more than 130 known interstellar and circumstellar molecules, and many more chains are likely to be detected in space once their laboratory spectra have been found and measured. New, highly precise measurements in the infrared and far-infrared region are very timely for the upcoming new generation of astrophysical instrumentation such as the Herschel satellite and the airborne observatory SOFIA (Stratospheric Observatory For Infrared Astronomy), covering the frequency domain in which many of the carbon chain spectra are presumably located.

The application of Terahertz- and infrared-spectroscopy to supersonic molecular beams has overcome the production difficulty, yielding the detection of various reactive and highly unsaturated carbon molecules. Recent results on pure carbon chain molecules and carbon hydrides will be presented and compared to high level *ab initio* calculations.

Molecular beam studies on reactions of C atoms and C₂ radicals with unsaturated hydrocarbons: a possible synthetic route to interstellar carbon chains

P. Casavecchia¹, F. Leonori¹, E. Segoloni¹, R. Petrucci¹, A. Bergeat², N. Balucani¹

¹Dipartimento di Chimica, Università di Perugia, 06123 Perugia, Italy

²LPCM, UMR5803, Université Bordeaux I, 33405 Talence Cedex, France

The reactions of ground-state carbon atoms, C(³P) and dicarbon molecules, C₂(X¹Σ_g⁺), with unsaturated hydrocarbons are of basic chemical interest and of great importance for understanding combustion processes and the chemical evolution of extraterrestrial environments such as molecular clouds and circumstellar envelopes of dying carbon stars, as for instance, IRC + 10216. Kinetic studies found these neutral-neutral reactions very fast down to very low temperature (15 K for C [1] and 24 K for C₂ [2]) and suggested that these are barrierless reactions that may have a critical role in the chemistry of the interstellar medium (ISM). An issue of key importance is how larger molecules are assembled and, in particular, how molecules containing chains of several carbon atoms are synthesized.

Here we report on the investigation of the dynamics of the prototypical and perhaps most important reaction of C and C₂, that with the acetylene molecule, by using the crossed molecular beam (CMB) scattering technique with mass-spectrometric detection [3]. Primary products and branching ratios were determined as a function of collision energy (i.e., relative translational energy). We exploit the capability (a) of generating *continuous* supersonic beams of C and C₂, (b) of crossing the reactant beams at variable angles (45°-90°-135°) to widen the range of collision energies, and (c) of detecting the products by “soft” electron-ionization [4,5].

Angular distributions and time-of-flight (TOF) spectra at $m/z = 37$ (C₃H) and 36 (C₃) were recorded in a wide range of collision energies ($E_c \sim 3.5$ -50 kJ/mol) for the reaction C(³P)+C₂H₂. We have identified the primary products of the competing pathways leading to *linear (l)/cyclic (c)*-C₃H + H and C₃ + H₂, derived their center-of-mass (CM) angular and translational energy distributions, determined the branching ratios, and characterized the reaction micro-mechanism [5]. Information on the *c*-C₃H/*l*-C₃H branching ratio and respective exoergicities are also obtained. The spin-forbidden pathway leading to C₃(X¹Σ_g⁺)+H₂(X¹Σ_g⁺) is rationalized in terms of intersystem crossing between triplet and singlet potential energy surfaces. Since our C beams contain, in addition to C(³P), also a significant concentration of excited C(¹D) atoms, we have characterized the dynamics of C₃H and C₃ formation from also the C(¹D)+C₂H₂ reaction (of relevance in comets). These studies extend significantly our previous investigation at $E_c = 29.3$ kJ/mol [6] and earlier CMB studies with pulsed beams [7]. Notably, from measurements of electron-ionization efficiency curves of the reaction products down to very low electron energies, the first experimental estimate of the ionization energy (IE) of *l*-,*c*-C₃H radicals is also obtained (the experimental IE compare well with theoretical predictions [8]). Angular distributions and TOF spectra at $m/z = 49$ (C₄H) from the reactions C₂(X¹Σ_g⁺/a³Π_u) + C₂H₂ were recorded at $E_c = 13.1$ and 36.8 kJ/mol. The product CM angular and translational energy distributions are determined. The dynamics of the buta-1,3-diynyl [C₄H(X²Σ⁺), HCCCC] radical product as a function of E_c is determined and compared with the results of recent CMB experiments carried out using pulsed beams [9].

References

- [1] D. Chastaing, S.D. Le Picard, I.R. Sims and I.W.M. Smith, *Astron. Astrophys.* **365**, 241 (2001).
- [2] I. R. Sims, A. Canosa, S. Le Picard *et al.*, Communication in *Faraday Discuss.* **133** (2006), in press.
- [3] P. Casavecchia, *Rep. Prog. Phys.* **63**, 355 (2000).
- [4] P. Casavecchia *et al.*, *J. Phys. Chem. A* **109**, 3527 (2005).
- [5] M. Costes, P. Casavecchia *et al.*, *Faraday Discuss.* **133** (2006), in press.
- [6] L. Cartechini *et al.*, *J. Chem. Phys.* **116**, 5603 (2002).
- [7] R. I. Kaiser, W. Sun, A. G. Suits, and Y. T. Lee, *J. Chem. Phys.* **107**, 8713 (1997).
- [8] R. K. Chaudhuri, S. Majumder, K. F. Freed, *J. Chem. Phys.* **112**, 9301 (2000).
- [9] R. I. Kaiser *et al.*, *Chem. Phys. Lett.* **382**, 112 (2003); *Faraday Discuss.* **133** (2006), in press.

Intra- and intermolecular dynamics of cyanoacetylene and its complexes with helium

Robert Moszynski¹, Leonid Shirkov¹, and Marie-Lise Dubernet²

¹Quantum Chemistry Laboratory, University of Warsaw, Poland

²Observatoire de Paris, Université de Paris VI, France

Accurate potential energy surfaces and nearly exact methods describing the nuclear motions are of primary importance to understand spectroscopic and collisional properties of molecules. In my talk I will show on the example of cyanoacetylene how modern quantum chemical methods can be applied with trust to describe state resolved spectroscopic and collisional experiments. Special attention will be paid to the state-of-the-art *ab initio* techniques for the calculation of the intramolecular potential energy surface of cyanoacetylene and to methods needed on the route from the potential energy surface to the rovibrational spectra. The exact Hamiltonian describing the nuclear motions will be introduced, and a hierarchy of possible decoupling schemes for an approximate calculation of the rovibrational energy levels will be discussed. Nearly exact energy levels from variational 10D calculations, as well as some approximate values obtained by using some decoupling procedures, will be reported and compared with high-resolution experimental data from microwave and infrared spectroscopy.

Polyynes And Cyanopolyynes: From Astrochemistry To Their Easy Synthesis In A Carbon Arc

Franco Cataldo

Soc. Lupi Chemical Research Institute, Via Casilina 1626/A, 00133 Rome, Italy

A parallelism has been discovered between the polyynes formed by laser ablation experiments of graphite targets and those produced from the submerged electric arc between graphite electrodes. In both cases the same products are obtained.

Hydrogen-terminated polyynes are obtained when the carbon arc is struck under water; monocyano-polyynes (mixed with hydrogen-terminated polyynes) are the products if the carbon arc is sparked in acetonitrile or ammonia; dicyano-polyynes if the arc is struck in liquid nitrogen. The mechanism of formation of polyynes in the submerged carbon arc involves essentially neutral species as it happens around the circumstellar environment.

It is shown that the relative abundances of the polyynes formed in the submerged carbon arc or in a carbon arc in vacuum decreases by a factor between 3 to 5 as the chain length increases by a C_2 unit. Exactly the same trend has been observed both for polyynes and cyanopolyynes in the circumstellar environment around carbon-rich stars. This fact suggests that the mechanism of formation of the polyynes in the carbon arc is the same occurring in the surroundings of the carbon-rich stars.

References

- [1] F. Cataldo (editor) *Polyynes: Synthesis, Properties and Applications*. CRC Press (Taylor & Francis Publishing Group), Boca Raton, FL, USA, 2005.
- [2] F. Cataldo. Cyanopolyynes: carbon chains formation in a carbon arc mimicking the formation of carbon chains in the circumstellar medium. *International Journal of Astrobiology* **3** (2004) 237-246.
- [3] F. Cataldo. Monocyano-polyynes from carbon arc in ammonia. About the relative abundance of polyynes series formed in a carbon arc and those detected in circumstellar medium. *International Journal of Astrobiology*. **5** (2006), in press.

Formation of interstellar methanol and other oxygen containing compounds: Non-feasibility of gas-phase mechanisms

W. D. Geppert¹, M. Hamberg,¹ R. D. Thomas¹, F. Österdahl², F. Hellberg¹, V. Zhaunerchyk¹, A. Ehlerding¹, T. J. Millar³, H. Roberts³, J. Semaniak⁴, M. af Ugglas¹, A. Källberg⁵, A. Simonsson⁵, M. Kaminska⁴ and M. Larsson¹.

¹ Molecular Physics Division, Department of Physics, Stockholm, Sweden.



² Royal Institute of Technology, AlbaNova, SE 10691, Stockholm, Sweden.

³ School of Physics and Astronomy, University of Manchester, Sackville St Building, PO Box 88, Manchester M60 1QD, UK

⁴ Institute of Physics, Świętokrzyska Academy, PL-25406, Kielce, Poland

⁵ Manne Siegbahn Laboratory, Frescativägen 24, SE-104 05 Stockholm, Sweden

For astronomical observers, methanol is one of the most interesting molecules. Multi-line observation on methanol can be used to determine temperature and spacial density of in an interstellar cloud with *one* molecule only, thereby avoiding the problem of different spatial distributions of different tracers.² Masers of methanol are seen as excellent tracers of star formation.³ Furthermore, the compound can be used as an evolution indicator during the embedded phase of massive star formation.⁴ Contrary to its importance for astronomical observations, comparatively little information is available on the formation of interstellar methanol. A two-step mechanism including radiative association of CH₃⁺ and H₂O and dissociative recombination of the produced ion has hitherto been held responsible:



To assess the feasibility of this process in the gas-phase the efficiency of reactions 1 and 2 has to be assessed. Therefore, branching ratios of the different reaction pathways and the overall rate coefficients of the dissociative recombination of CH₃OH₂⁺ (Reaction 2) has been measured at the CRYRING storage ring located in Stockholm, Sweden. Analysis of the data yielded that formation of methanol accounted for only 3 % of the total rate in CH₃OH₂⁺ (6 % in the isotopomer CD₃OD₂⁺). Largely, dissociative recombination of CH₃OH₂⁺ involves fragmentation of the C-O bond, the major process being the three-body break-up forming CH₃, OH and. The overall cross section is best fitted by $\sigma = 1.2 \pm 0.1 \times 10^{-15} E^{-1.15 \pm 0.02} \text{ cm}^2$. From these value the thermal reaction rate coefficients of $k(T) = 8.9 \pm 0.9 \times 10^{-7} (T/300)^{-0.59 \pm 0.02} \text{ cm}^3 \text{ s}^{-1}$ can be calculated.

These values, along with a recently measured upper limit for the rate coefficient of reaction (1)⁵ have been used as input for a model calculation of a dark interstellar cloud resembling TMC-1 using the UMIST code.⁶ This calculation yields a peak methanol abundance of 7×10^{-13} (H₂) which is definitely below the observed values in dark clouds and probably below the detection limits for existing telescopes. A non-negligible formation of interstellar methanol by the previously proposed mechanism is therefore very unlikely and a grain surface process for methanol production *via* repetitive hydrogenation of CO is invoked. Very recently, it has been established by Hidaka *et al.*⁷ in a surface reaction experiment that the formation of methanol by hydrogenation of CO with H on pure CO and mixed CO/H₂O ices proceeds efficiently at 10 K. Moreover, once produced, CH₃OH is not degraded by irradiation.

Very recently, the dissociative recombination of protonated dimethylether (CH₃)₂OH⁺ has also been investigated, which has been regarded as the final step in formation of interstellar dimethyl ether.⁸ Preliminary analysis shows that the branching ratio for dimethyl ether formation in this reaction amounts to only 6 %, which, in analogy to the findings on methanol, renders the previously assumed formation pathway very improbable.

² S. Leurini, P. Schilke, K. M. Menten, D. R. Flower, J. T. Pottage, L. H. Xu, *Astron. Astrophys.* 2004, **422**, 573

³ V. Minier, *Technical Report No. 395, Chalmers University of Technology, Gothenburg*, 2000

⁴ F. F. S. van der Tak, E. F. van Dishoeck, P. Caselli, *Astron. Astrophys.*, 2000, **361**, 327

⁵ A. Luca, D. Voulot, D. Gerlich, *WDS'02 Proceedings of Contributed Papers, PART II*, 2002, 294

⁶ A. Markwick, T. J. Millar, S. B. Charnley, 2000, *Astrophys. J.*, **535**, 256

⁷ H. Hidaka, N. Watanabe, T. Shiraki, A. Nagaoka, A. Kouchi, 2004, *Astrophys. J.*, **614**, 1124.

⁸ S. B. Charnley, M. E. Kress, A. G. G. M. Tielens, T. J. Millar, *Astrophys. J.* **448**, 232

Experimental studies of H₂ formation on grains: overview and recent results

Francois Dulieu

LERMA, Université de Cergy Pontoise, 5 Mail Ga-Lussac, Cergy-Pontoise, 95031 France

Irradiation and thermal processing of silicates in the laboratory: Insights in extraterrestrial grain properties

Z. Djouadi¹, C. Davoisne², L. d'Hendecourt¹, H. Leroux², A. Jones¹

¹ Institut d'Astrophysique Spatiale (IAS), bâtiment 121, F-91405 Orsay, Université Paris-sud 11-CNRS (UMR 8617) France.

² Laboratoire de Structure et Propriétés de l'Etat Solide (LSPES)-UMR 8008, Université des Sciences et Technologies de Lille, 59655 Villeneuve d'Ascq Cedex, France.

Before the results from the Short Wavelength Spectrometer instrument (SWS) on board the Infrared Space Observatory (ISO), it was largely thought that all silicate dust in space was totally amorphous. The extensive results provided by ISO have shown the existence of different forms of dust (crystalline, amorphous and/or both of them), depending essentially on the astrophysical environment where they are detected. Many studies have been devoted to understand the primary modifications (structure and chemical composition) due to ionic irradiations [1-3] or thermal processing [4-5], but very little has been done regarding further evolution of irradiated silicates. In order to evaluate the dust ability to crystallize in cold environments, we have studied thermal processing of silicates after a further irradiation stage [6]. For this, we have developed an experimental protocol simulating different stages of the life-cycle of the dust. Thin silicate films (typically 50 to 100 nm in thickness) are synthesized by electron-beam evaporation of San-Carlos olivine ($\text{Mg}_{1.8}\text{Fe}_{0.19}\text{Ni}_{0.01}\text{SiO}_4$) onto 3 mm diameter diamond discs. He^+ irradiations, under vacuum at room temperature at low energies (5 keV and 10 keV) and at fluences similar to those of interstellar shock waves (10^{16} and 10^{17} cm^{-2}) were used in order to completely amorphize the samples. Thermal processing was performed in a tubular furnace under vacuum at different temperatures. The evolution of the silicate structure was monitored by infrared spectroscopy. Using the 10 μm band evolution with annealing time we have determined the activation energy of crystallization after an irradiation stage [6]. Our result clearly suggests that the presence of crystalline silicates in cold environments can not be explained by a local thermal diffusion process. One can thus invoke the radial mixing between the inner regions of the nebula and the outer cooler and comet-forming regions [7].

The smooth surface of the diamond allows an easy extraction of the film for TEM characterization. Microstructural analyses are obtained with a Philips CM30 TEM operating at 300 kV and the crystallographic characterization is achieved using selected area electron diffraction (SAED). The microscope is also equipped with an X-ray energy dispersive spectrometer (EDS) for microanalysis. We show the possibility to form spheroid metallic (iron and nickel) nano-particles (2-50 nm) randomly distributed within the silicate films, which are still amorphous after annealing at 870 K even for long time duration (up to 780 hours). The iron initially present in form of FeO has thus segregated from the amorphous phase in the form of metallic globules (Figure1). This clearly suggests that a reduction has occurred according to the reaction $\text{FeO} \rightarrow \text{Fe} + \frac{1}{2} \text{O}_2$. The obtained microstructures closely resemble those of the GEMS (Glass Embedded with Metal and Sulfides) found in chondritic IDPs (Interplanetary Dust Particles) [8]. Locking iron as metallic particles within the silicates explains first why astronomical silicates always appear observationally Fe-poor. Secondly, these metallic inclusions could explain the dust alignment in the galactic magnetic field. Measurements to determine the magnetic susceptibility of our minute samples are under progress with the goal of understanding the magnetic properties of such small metallic iron inclusions in interstellar grains.




Figure1: TEM micrograph of the sample before (on the left) and after (on the right) annealing at 600°C during ~ 800 hours. Note the presence of inclusions in the heated sample which is still amorphous as showed by the diffraction pattern given on the top right.

References: [1] Demyk K., et al. (2004) *A&A*, 420, 233. [2] Carrez P., et al. (2002), *Meteor. Planet. Sci.* 37, 1599. [3] Brucato J.R., et al. (2004), *A&A*, 413, 395. [4] Brucato J.R., et al. (2002), *PSS*, 50, 829. [5] Hallenbeck S.L. et al. (1998), *Icarus*, 131, 198. [6] Djouadi Z., et al. (2005) *A&A*, 179-184. [7] Bockelée-Morvan et al, (2002), *A&A*, 384, 1107. [8] Davoisne C. et al, (2006) *A&A*, 448, L1.

Photo- And Radoracemization Of Chiral Molecules And The Radiation-Induced Amplification Of Chirality

Franco Cataldo

Soc. Lupi Chemical Research Institute, Via Casilina 1626/A, 00133 Rome, Italy

The biochemistry of the biological world is based on molecules and macromolecules having one or more centers of asymmetry, these molecules are called chiral molecules. Thus, life is strictly linked to the chirality of the molecules and macromolecules. No life can be conceived without chirality because all the complex chemistry and biochemistry, i.e. high selectivity in chemical reactions in living organism, which are the basis of life would be impossible.

A central problem of the research of the origin of life is the origin of chirality. Phenomena like the action of circularly polarized light on racemic mixture of certain molecules have been pointed out as the spontaneous source of chiral excess in an abiotic world. Thus, it is thought that the chiral molecules are formed already in the interstellar medium, in the molecular clouds and then they are incorporated into primitive bodies like comets and meteorites. Experimental evidences are of support to these hypotheses.

However, there are two phenomena which play against the preservation of chirality of asymmetric molecules in space: the photoracemization and the radoracemization. In fact the photolysis and the radiolysis of chiral molecules caused by high energy radiation and by cosmic rays in general cause a reduction in the optical activity. Apparent and true photo- and radoracemization must be distinguished.

An additional problem must be discussed, the fact that the chiral excess produced by spontaneous phenomena like the action of circularly polarized light on racemic mixtures yields only scalemic mixtures with a small enantiomeric excess which is further penalized by the natural photo- and radoracemization phenomena. Therefore, natural amplification mechanisms of chirality should exist in order to be able to increase the enantiomeric excess of scalemic mixtures and make them useful for the emergence of the primordial life.

Very recently we have demonstrated that such mechanism of chiral amplification is offered for examples by the radiopolymerization of chiral monomers which are able to yield products with higher optical activity than the starting monomer.

References

- [1] F. Cataldo, Y. Keheyan. Radiopolymerization of β (-)-pinene: a case of chiral amplification. *Radiation Physics and Chemistry*, 75 (2006) 572-582.
- [2] F. Cataldo, J.R. Brucato, Y. Keheyan. Chirality in prebiotic molecules and the phenomenon of photo- and radoracemization. *Journal of Physics Conference Series*, 6 (2005) 139-148.
- [3] F. Cataldo, Y. Keheyan and S. Baccaro. Gamma-radiolysis of chiral molecules: R(+)-limonene, S(-)-limonene and R(-)-alpha-phellandrene. *Journal Radioanalytical and Nuclear Chemistry*, 262 (2004) 423-428.
- [4] F. Cataldo, J.R. Brucato, Y. Keheyan. Gamma radiation induced polymerization of a chiral monomer: a new way to produce chiral amplification. *Origin of Life and Evolution of Biosphere*, in press.
- [5] F. Cataldo, Y. Keheyan. An essay on the asymmetric polymerization and the origin of chirality in biological macromolecules. *Journal of Chemical and Biochemical Kinetics* 5 (2003) 135-155. Also published in the book: G.E. Zaikov (editor) International Perspectives in Chemistry and Biochemistry Research. Nova Science Publishers, New York, Chapter 3, (2003)

Spectroscopy of complex and organic chemistry in Titan.

A. Jolly

LISA, University Paris 12, 61 Avenue du General de Gaulle, Paris 94010, France

List of Posters

- P1 Can amino acids be formed during the evolution of molecular cloud?**
Kinsuk Acharyya, Sandip K. Chakrabarti and Sonali Chakrabarti
- P2 Detection of the gaseous disk around the B0 star R Mon**
T. AlonsoAlbi, A. Fuente, R. Bachiller, A. Natta, L. Testi, R. Neri, P. Planesas
- P3 The Far-Infrared Fourier Transform Emission Spectrum of HCN**
D. Bailly, O. Pirali, M. Vervloet
- P4 Complex reaction networks on dust grains in dense molecular clouds**
Ofer Biham, Baruch Barzel and Azi Lipshtat*
- P5 A 80 - 370 GHz Unbiased Spectral Survey of the Solar-Type protostar IRAS 16293-2422**
E. Caux, A. Bacmann, S. Bisschop, S. Botinelli, A. Castets, S. Cazaux, C. Ceccarelli, C. Comito, F.P. Helmich, C. Kahane, B. Parise, P. Schilke, A.G.G.M. Tielens, E.F. van Dishoeck, V. Wakelam, A. Walters
- P6 Monte Carlo Simulation of the Production of Hydrogen Molecules on Grain Surfaces**
Sandip K. Chakrabarti, Ankan Das, Kinsuk Acharyya, Sonali Chakrabarti
- P7 Time evolution of the Complex molecule formation in collapsing interstellar cloud**
Sandip K. Chakrabarti, A. Das, K. Acharyya and S. Chakrabarti
- P8 Constraints on dark clouds structures based on molecular lines observations**
Fabien Daniel, Observatoire de Paris
- P9 Time dependent chemical evolution of molecular cloud**
Ankan Das, S.K. Chakrabarti, Kinsuk Acharyya, S. Chakrabarti
- P10 Disentangling the structure of intermediate mass hot cores at scales of ~200 AU**
A. Fuente, C. Ceccarelli, R. Neri, D. Johnstone, P. Caselli, F. Wyrowski, E. van Dishoeck
- P11 Dissociative Recombination of sulfur-containing interstellar ions**
Wolf Dietrich Geppert, Stockholm University
- P12 Terahertz Laboratory Spectroscopy**
Thomas Giesen, University of Cologne
- P13 Turbulence, star formation and the need for high spatial resolution**
M. Gustafsson, A. Brandenburg, J.L. Lemaire, F.P. Pijpers, D. Field
- P14 Laboratory simulations of molecular hydrogen formation in PDRs**
Liv Hornekaer, University of Aarhus

- P15 Search for PAH emission features in the far-IR/submm range**
Christine Joblin, CNRS
- P16 Clumping and photochemistry in planetary nebulae**
Eric Josselin, Université Montpellier II
- P17 Modelling water in the envelopes of low-mass protostars**
T.A. van Kempen, S. D. Doty, E.F. van Dishoeck, M.R. Hogerheijde, J.K. Jørgensen
- P18 Deuterated chemistry on amorphous water ice surfaces in the ISM**
L. Kristensen, L. Amiaud, F. Dulieu, J-H. Fillion, A. Momeni, J. L. Lemaire
- P19 Evaluation of spectral data for radioastronomy, Herschel and ALMA. Contribution to the improvement of this data.**
Valerio Lattanzi, Brian Drouin, John Pearson, Adam Walters
- P20 Formation and detection of complex molecules in astrochemical ice**
Nigel Mason, The Open University
- P21 Astrochemistry with the Atacama Pathfinder Experiment**
Karl Menten, Max-Planck Institute for Radioastronomy
- P22 Velocities of gas in star-forming regions: 70AU resolution and 1km/s results for OMC1.**
H.D.Nissen, M.Gustafsson, J.L.Lemaire, Y.Clénet, D.Rouan, D.Field
- P23 Alcohol Chemistry In The Galactic Center. Hot Core Chemistry Without Hot Cores.**
M. A. Requena-Torres, J. Martín-Pintado, A. Rodríguez-Franco, S. Martín, N. J. Rodríguez-Fernández and P. de Vicente
- P24 The survival of dense gas around pre-WR stars**
J. Ricardo Rizzo, Universidad Europea de Madrid
- P25 Low-mass starformation in the R CrA molecular cloud**
Fredrik Schöier, Stockholm University
- P26 Radiative lifetimes and association cross-sections of CO²⁺**
T. Šedivcová, P. R. Ždánková and V. Špirko
- P27 Reliability of the disk chemical modeling**
A.I. Vasyunin, D. Semenov, A.M. Sobolev, Th. Henning
- P28 Rotational excitation of interstellar cyanoacetylene**
Wernli M, Wiesenfeld L, Faure A, Valiron P
- P29 Planet Formation - Traced with ALMA**
Sebastian Wolf, Dmitry Semenov, Gennaro D'Angelo, Amaya Moro-Martín, and Yaroslav Pavlyuchenkov

Can amino acids be formed during the evolution of molecular cloud?

Kinsuk Acharyya¹, Sandip K. Chakrabarti^{1,2} and Sonali Chakrabarti^{1,3}

¹Centre for space Physics, 43 chalantika, Garia, Kolkata, 700084,

²S.N. Bose National Centre for Basic Sciences, JD-Block, Sector III, Salt Lake,
Kolkata 700091, India and

³Maharaja Manindra Chandra College, Kolkata

Using a reasonably large gas phase chemical reaction network consisting of 421 species and H₂ formation on grain we investigate the formation of simple amino acids like, glycine and alanine. We studied the chemical evolution of molecular clouds during the static and the collapse phase of star formation. These findings may shed light on the present debate on glycine detection in molecular clouds. We found glycine could form in detectable amount in both static and collapsing phase of the molecular cloud. However, we find that β -alanine and propionic acid is not produced in detectable amount in both types of clouds.

Keywords: ISM: molecules, Molecular clouds, Star formation, Chemical evolution, Bio-molecules

Detection of the gaseous disk around the B0 star R Mon

T. AlonsoAlbi¹, A. Fuente¹, R. Bachiller¹, A. Natta², L. Testi², R. Neri³,
P. Planesas¹


¹ Observatorio Astronómico Nacional, Apdo. 112, E28803 Alcalá de Henares, Spain

² Observatorio Astrofisico di Arcetri, INAF, Largo E. Fermi 5, 50125 Firenze, Italy

³ Institut de Radio Astronomie Milimétrique, 300 Rue de la Piscine,
Domaine Universitaire de Grenoble, F38406 St. Martin d'Hères, France

We present high angular resolution observations of the ^{12}CO 1 \rightarrow 0 (beam = 2.3'' x 1.3'') and 2 \rightarrow 1 (beam = 1.5'' x 0.9'') rotational lines. At this scale (1'' = 800 AU at the distance of R Mon) the high velocity CO emission has two components: the high velocity gas flowing outwards the star along twisted filaments located in the walls of the nebula cavity, and the circumstellar disk. We have used a plane disk model to fit the ^{12}CO 1 \rightarrow 0 and 2 \rightarrow 1 emission in a strip perpendicular to the outflow axis and to determine the kinematics of the gaseous disk. The model assumes standard ^{12}CO abundance ($X(\text{CO}) = 8 \cdot 10^5$), radial dependent temperature and density laws (T (K) = $T_0 r^{-\alpha}$, ρ (cm $^{-3}$) = $\rho_0 r^{-\beta}$), and Local Thermodynamic Equilibrium. The CO emission is consistent with a plane disk at the inclination angle of 20° in keplerian rotation around the star. The disk is fitted with a mass of 0.01 M_{\odot} , an outer radius of 1500 AU, a temperature of 4000 K at the inner radius of 1 UA, and values of $\alpha = 0.65$ and $\beta = 1.3$ for the indexes of the temperature and density laws.

In addition, we present new continuum observations of the region with a beam $\sim 0.72'' \times 0.33''$. These observations provide more accurate estimates of the position and size of the dusty disk, which are now settled to RA (2000): 06:39:09.96, DEC(2000): 08:44:09.55, and rout # 150 UA.



Right: Interferometric map in the ^{12}CO 2-1 line of the red ($v < 7$ km/s) and blue ($v > 12$ km/s) lobes of the outflow associated with the B0 star R Mon. A strip perpendicular to the outflow is drawn in this panel. Left: PositionVelocity (P-V) diagrams of the emission of the ^{12}CO 1-0 and 2-1 lines along the strip drawn in the top panel. These P-V diagrams are consistent with the existence of a disk in keplerian rotation around the star. We have overlaid in red contours the results obtained with our disk model. Note that the emission at the velocities of the molecular cloud (9.5 ± 2.5 km/s) has been filtered out by the interferometric observations.

The Far-Infrared Fourier Transform Emission Spectrum of HCN

D. Bailly¹, O. Pirali², M. Vervloet¹

¹ LPPM, CNRS, Bât 350, Université Paris-Sud, Orsay, 91405, France

² FOM Institute for Plasma Physics, Edisonbaan 14, Nieuwegein, 3439MN, The Netherlands

The rotational emission spectrum of hydrogen cyanide has been recorded by means of a high resolution Fourier transform spectrometer. Excited HCN was created in a low pressure N₂/C₂H₄ mixture flowing through a radio-frequency discharge. The spectrum covers the far-infrared spectral range 26-180 cm⁻¹ (0.8-5.4 THz). No lines of HNC have been clearly identified. All the rotational lines of HCN assigned in the spectrum belong to its 15 lowest vibrational states. The highest observed state, namely (050), has been studied for the first time by rotational spectroscopy. It is perturbed by (021) through Coriolis interaction. Three vibration-rotation lines of the band (050) – (021) are identified in the spectrum in the range 76-80 cm⁻¹ where the Coriolis interaction reach a maximum for J = 26. These lines have been found to produce laser emissions from electric discharges in compounds of hydrogen, carbon and nitrogen (1). A similar situation occurs at lower frequency (at about 30 cm⁻¹) between (040) and (011) and has been studied in a previous submillimeter-wave spectroscopy work (2). Strong maser emission have been detected between (040) and (011) at 805 and 891 GHz by Schilke et al. (3) and Schilke and Menten (4) respectively.

We have deduced molecular constants for the 15 lowest vibrational levels of HCN by combining our measurements with those reported in (2). They show a very good agreement with the results obtained from high resolution infrared emission spectroscopy (5). We hope they will help in the identification of highly excited HCN in the far infrared region.

1 L. E. S. Mathias, A. Crocker, and M. S. Wills, IEEE J. Quantum Electronics, vol. QE-4 (1968) 205-208.

2 Z. Zelinger, T. Amano, V. Ahrens, S. Brünken, F. Lewen, H. P. S. Müller, and G. Winnewisser, J. Mol. Spectrosc., 220, (2003) 223-233.

3 P. Schilke, D. M. Mehringer, K. M. Menten, Astrophys. J., 528 (2000) L37- L40.

4 P. Schilke, K. M. Menten, Astrophys. J., 583 (2003) 446-450.

5 A. G. Maki, G. Ch. Mellau, S. Klee, M. Winnewisser, and W. Quapp, J. Mol. Spectrosc., 202 (2000) 67-82.

Complex reaction networks on dust grains in dense molecular clouds

Ofer Biham, Baruch Barzel and Azi Lipshtat*

Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel


There are strong indications that networks of chemical reactions on dust grains play an important role in the production of molecules such as methanol and its deuterated versions in molecular clouds [1]. Due to the small size of the grains and the low flux, the populations of adsorbed atomic and molecular species are small and exhibit large fluctuations. As a result, rate equations are not suitable for the simulation of reaction networks on grains, which requires stochastic methods such as Monte Carlo simulations or direct integration of the master equation. An important advantage of the master equation is that it consists of differential equations which can be easily coupled to the rate equations of interstellar gas-phase chemistry. However, the number of equations proliferates making the master equation infeasible for complex reaction networks. Here we present two methods which provide a dramatic reduction in the number of equations. The multi-plane method maintains the structure of the master equation, but dramatically reduces the number of equations [2]. In the second method, based on moment equations, the number of equations is further reduced making it feasible to incorporate stochastic modelling of grain-surface reaction networks into models of interstellar chemistry.

[1] T. Stantcheva and E. Herbst, MNRAS 340, 983 (2003); R. Garrod, I.H Park, P. Caselli and E. Herbst, Faraday Discuss. (2006).

[2] A. Lipshtat and O. Biham, Phys. Rev. Lett. 93, 170601 (2004).

* Present address: Dept. of Pharmacology and Biological Chemistry, Mount Sinai School of Medicine, New York, NY 10029, USA

A graph describing the surface reaction network that produces methanol. The reactive species are shown in the nodes. Species that react with each other are connected by an edge. The reaction products are specified near the edges.



A 80 - 370 GHz Unbiased Spectral Survey of the Solar-Type protostar IRAS 16293-2422

E. Caux¹, A. Bacmann², S. Bisschop³, S. Botinelli⁴, A. Castets², S. Cazaux⁵,
C. Ceccarelli⁴, C. Comito⁶, F.P. Helmich⁷, C. Kahane⁴, B. Parise^{6,1}, P.
Schilke⁶, A.G.G.M. Tielens⁸, E.F. van Dishoeck³, V. Wakelam^{9,2}, A. Walters¹

¹CESR CNRS-UPS, BP 4346, 31028 Toulouse cedex 04, F

²Observatoire de Bordeaux, BP 89, 33270 Floirac, F

³Leiden Observatory, PO Box 9513, 2300 RA Leiden, NL

⁴LAOG, BP 53, F-38041 Grenoble cedex 09, F

⁵INAF, Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, 50125 Firenze, I

⁶Max-Planck Institute für RadioAstronomie, Auf dem Hugel 69, 53121 Bonn, G

⁷SRON, PO Box 800, 9700 AV Gröningen, NL

⁸Kapteyn Astronomical Institute, P.O. Box 800, 9700 AV Groningen, NL

⁹The Ohio State University, Woodruff Ave. Columbus, OH 43210-1106, USA

Unbiased line surveys are a unique way to determine the molecular content of astrophysical objects. They allow a complete study of the physics and chemistry in and around molecular sources. Among all class “0” solar-type protostars observed to date, IRAS16293-2422 has the richest line spectrum. It is the first one where several complex molecules have been observed (Cazaux et al. 2003) as well as many deuterated molecules (D₂CO, Ceccarelli et al. 1998 and doubly and triply deuterated methanol, Parise et al. 2002, 2004). Ceccarelli et al. 1998 and Shoier et al. 2002 have shown that the structure of the IRAS16293 collapsing envelope has a inner hot corino ($r < 150$ AU) where the temperature exceeds 100 K and a colder outer envelope. This spatial structure is responsible for the abundance jumps for several molecules.

Due to its high temperature and density, this hot corino is responsible for the richness of the IRAS16293 spectrum. This richness comes also from the molecular content of the source itself, due both to the release of molecules trapped in icy mantles by the elevated temperatures, and to gas phase chemical reactions which can proceed faster at high density. High temperatures open many chemical pathways which, because they are endothermic, are closed at lower temperatures. We present here preliminary results of this 80 - 370 GHz unbiased spectral line survey.

IRAS16293 has a very rich molecular spectrum (~20 lines / GHz), regardless of the frequency. The total number of lines expected is a few thousand that constrained us to develop a specific software to handle these data (CASSIS, see the poster from A. Walters et al., this conference), One can note the presence of many lines originating from various molecules including complex and deuterated molecules. The line identification of the whole survey is in progress, and we plan to make public the line database by April 2007.

This kind of broad and unbiased survey is mandatory to obtain acceptable accuracy on the physical parameters. For most species, we will first derive the physical parameters (Tex and column density) from rotational diagrams. Species for which collisional coefficients are known will be analysed using more sophisticated models (LVG...), as done for the emission of HDO in IRAS16293 from this survey (Parise et al. 2005). A complete census of the molecular content of IRAS16293 will be built, and the detailed modelling of the chemistry governing its hot core and envelope will allow the comparison with massive hot core regions and will put stringent constraints on chemical models.

Monte Carlo Simulation of the Production of Hydrogen Molecules on Grain Surfaces

Sandip K. Chakrabarti^{1,2}, Ankan Das², Kinsuk Acharyya²,
Sonal Chakrabarti^{2,3}

¹S.N. Bose National Centre for Basic Sciences, JD Block, Salt Lake, Kolkata,
700098, India;

²Centre for Space Physics, 43 Chalantika, Garia Station Rd., Kolkata, 700084, India;

³Maharaja Manindra Chandra College, Ramakanta Bose Street, Kolkata 700003, India

Majority of the observed hydrogen molecules in the interstellar medium are formed on the grain surfaces. Dust grains are essential for the formation of the hydrogen molecule. On the grain surfaces, two H atoms combine to form a single hydrogen molecule. So far, the average recombination time to produce a hydrogen molecule used to be computed by a crude assumption about the diffusion of the H atom on a grain surface. Here, we perform a Monte-Carlo simulation to exactly quantify how the diffusion actually takes place. This enables us to calculate the average recombination time for the formation of hydrogen molecules. We find that the production rate of the hydrogen molecules on the grain surfaces is highly dependent on the grain size and the accretion rate. This is a very important conclusion in the context of the grain chemistry and it will affect the molecular abundances of hydrogen, and many complex molecules in the evolving cloud. We perform our calculations on two types of grain surfaces namely, on the amorphous carbon and the Olivine grain surfaces for a wide variety of grain parameters.

Time evolution of the Complex molecule formation in collapsing interstellar cloud

Sandip K. Chakrabarti, A. Das, K. Acharyya and S. Chakrabarti

S.N. Bose National Centre for Basic Sciences (AA) Centre for Space Physics (AA,AB, AC, AD)

We study the time evolution of the interstellar cloud and find how the density and velocity profiles evolve. With this hydrodynamics, we couple chemical evolution consisting of a large network of species and reaction cross-sections. For molecular hydrogen formation we utilise the grain chemistry by master equation or rate equation depending on the accretion rates. We also present Monte-Carlo simulation to show that the way the H₂ molecular hydrogen formation is handled in the literature is seriously flawed. This is because we find that the effective recombination time of atomic hydrogens depend on the relative size of the grain and the accretion rate of the atomic hydrogen. We show how the chemical composition of the cloud evolves with time for various cloud and grain parameters.

Time dependent chemical evolution of molecular cloud

Ankan Das¹, S.K. Chakrabarti^{2,1}, Kinsuk Acharyya¹, S. Chakrabarti^{1,3}

¹Centre for Space Physics, 43 Chalantika, Garia Station Rd., Kolkata, 700084, India;

²S.N. Bose National Centre for Basic Sciences, JD Block, Salt Lake, Kolkata, 700098, India;

³Maharaja Manindra Chandra College, 20 Ramkanta Bose Street, Kolkata 700003

More than 125 species of molecules have been observed in the interstellar clouds and star forming regions. Among them over half are organic. In the present work, we try to explain the observational abundances. We develop a time-dependent hydro-dynamic code which determines the hydrodynamic evolution of the molecular cloud. Using the output of this code, we study the chemical evolution of this molecular cloud with time. In the chemical evolution, we incorporate the grain chemistry to accurately determine the abundance of H₂. We compare our results with observations.

Disentangling the structure of intermediate mass hot cores at scales of ~ 200 AU

A. Fuente¹, C. Ceccarelli², R. Neri³, D. Johnstone⁴, P. Caselli⁵, F. Wyrowski⁶, E. van Dishoeck⁷

¹Observatorio Astronómico Nacional, Apdo. 112, E28803 Alcalá de Henares, Spain

²LAOG, BP 53, F38041 Grenoble Cedex 9, France

³IRAM, 300 Rue de la Piscine, Domaine Universitaire, F38406 St. Martin d'Hères, France


⁴NRCC, Herzberg Institute of Astrophysics, 5071 West Saanich Rd, Victoria, BC, V9E 2E7, Canada

⁵INAF Osservatorio Astrofisico di Arcetri, Largo E. Fermi, 50125 Firenze, Italy

⁶MaxPlanck Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany

⁷Raymond and Beverly Sackler Laboratory for Astrophysics at Leiden Observatory, Postbus 9513, 2300 RA Leiden, The Netherlands

Hot cores are objects characterized by warm temperatures ($T > 100$ K), high density ($n > 10^6$ cm⁻³), and a very rich chemistry in complex molecules (CH₃OH, CH₃CN, CH₃CHOOH, CH₃OCH₃ ...). Hot cores have been thought to be associated with highmass protostars ($M > 10 M_{\odot}$) and represent an important phase in their evolution towards ultracompact and compact HII regions. Only recently, regions similar to hot cores have been detected in lowmass protostars. However, the amount of warm material involved, as well as the chemistry, are different in the two classes of objects and rise questions on the mechanisms that lead to the observed chemical complexity (Cazaux et al. 2003, ApJL 593, L51; Fuente et al., 2005, A&A 444, 481). The study of hot cores in intermediate mass protostars ($M_{*} \sim 2-10 M_{\odot}$) will provide an important clue to solve this problem. We have successfully searched for intermediate mass (IM) hot cores using the IRAM 30m telescope. In addition we have mapped four of these objects, IC1396 N, OMC2FIR4, CB 3, and Serpens-FIRS 1, using the new A+ configuration of the Plateau de Bure Interferometer. This configuration provides an angular resolution of $\sim 0.2''$ which implies a spatial resolution of < 200 AU at the distance of these IM hot cores. This high angular resolution allows us to measure with unprecedented accuracy the position, sizes and spectral indexes of these hot cores. Furthermore, these observations reveal the existence of a "cluster" of hot cores in the interior of IC 1396 N and CB 3. Simultaneously with the continuum observations we have mapped the region in the CH₃CN 13k \rightarrow 12k line and some complex Obearing molecules. Since the interferometric observations resolve for the first time the different components of the cluster, we are able to have a first glance at the chemical differentiation between them.



Continuum image at 1 mm of the Class 0 protostar IC 1396 N. A zoom in the central region is shown in the inset. Note that there is a cluster of 3 hot cores in the interior of this protostar. The spectrum towards the most massive component of the cluster is also shown.

Turbulence, star formation and the need for high spatial resolution

M. Gustafsson¹, A. Brandenburg², J.L. Lemaire³, F.P.Pijpers⁴, D. Field¹

¹ Dept. Physics and Astronomy, University of Aarhus, 8000 Aarhus, Denmark

² NORDITA, Blegdamsvej 17, 2100 Copenhagen Ø, Denmark

³ Observatoire de Paris & Université de Cergy-Pontoise, LERMA & UMR 8112 du CNRS, 92195 Meudon, France

⁴ Space and Atmospheric Physics, Dept. Physics, Imperial College London, England

Theories about the influence of turbulence on star formation need to be constrained by observations. Important issues such as the driving mechanism, the effect of re-injection of energy by protostellar outflows, the effect of self-gravity and the dissipation mechanism can only be settled by observations and simulations ability to reproduce the observations.

In order to assess the dissipation scale, conventionally the scale at which bulk energy is converted into heat, as well as the scales of low mass star formation it is essential to obtain maps with as high spatial resolution as possible.

We use infrared observations of shock-excited H₂ at 2.12 micron to characterise the velocity field in the star-forming region OMC1 at scales from 70 AU to 30000 AU. The results generally follow the trends observed at larger scales from radio observations tracing the cold, bulk gas. Hereby the validity of the ‘‘Larson Law’’ is extended by two orders of magnitude in scale. However the structure functions show the presence of preferred scales in the medium. These scales can possibly be associated with outflows from low mass protostars and are also the correct scale for disks around protostars. We compare the observations with subsets of simulations of (non-magnetic) hydrodynamical turbulence, which are forced at large scales and without self-gravity. To represent best the observations the selected subsets include only the shocked part of the simulations. The simulations are unable to reproduce the preferred scales found in the observations.

Performing structure analysis using data of warm H₂ has the intrinsic caveat that only shocked gas is observed. With ALMA it will be possible to map for example CO and trace the cold gas with as high or higher spatial resolution as in the exciting infrared data, thus making it possible to reach the dissipation scale of turbulence.

Modelling water in the envelopes of low-mass protostars

T.A. van Kempen¹, S. D. Doty², E.F. van Dishoeck¹, M.R. Hogerheijde¹,
J.K. Jørgensen³

¹Sterrewacht Leiden, Niels Bohrweg 2, 2311 CA, Leiden, Netherlands

²Dept. of Physics & Astronomy, Denison University, Olin 109 Granville, OH 43023

³Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

Water is one of the key molecules in the early stages of star formation, especially in the warm inner region of newly formed protostars, where ice mantles are evaporated from grains. In addition water is the basis of oxygen driven chemistry and essential for the creation of more complex molecules. The launch of the Herschel Space Observatory and the coming of ALMA will provide the opportunity to study gas-phase water in detail in YSOs at high sensitivity and resolution. The HIFI spectrometer on-board Herschel with a spectral resolution of $\lambda/\Delta\lambda = 10^7$ is especially suited to observe gas-phase rotational lines over a wide range of excitation conditions. ALMA provides the single most powerful probe into the warm inner region of circumstellar envelopes, which span ~tens of AU. Many isotopologues of water such as HDO and many complex molecules associated with water are also confined to these warm regions of the envelopes characteristic of Class 0 and Class I sources.

We present here the first results of a parameter study on the role of water in YSOs, focussing on low-mass Class 0 and Class I protostars (Lada 1987, André et al, 1993) using the radiative transfer code RATRAN (Hogerheijde & van der Tak, 2002). The study was done in collaboration with a similar study on high-mass stars (Doty et al, 2006 in prep). Several parameters were investigated that influence the behaviour of water. The luminosity, L , of a protostar, which depends on both age and initial mass, determines the local temperatures throughout the envelope and is thus responsible for the spatial extent of the warm inner core, where water is abundant due to evaporation of the grain mantles. Second, the density of the envelope varies between different young protostars (Jørgensen et al, 2002), both in the density at a set radius, n , as well as the radial density gradient, α . In general it is seen that older protostars, such as Class I have steeper density (~2) gradients, then Class 0 (~1.5) and larger warm inner regions. Third, not much is known about the abundance of water relative to H_2 , both in regions where water is abundant ($T > 100$ K) and where it is frozen out onto the grains ($T < 100$ K). See Fig. 1 for an overview of the structure of circumstellar envelopes.

The goal is to constrain H_2O abundances for different models and provide accurate models that ALMA can probe. Our first result show that many lines for water are observable for typical low-mass protostars, but that a thorough understanding of the physical characteristics of each individual protostar is essential in understanding any modelled or observed flux of a given water line. Figures 2 and 3 show the ground state (557 GHz) and the $2_{21}-2_{12}$ excited state (1661 GHz) lines of ortho-water. One can clearly see that for a given luminosity ($L = 2 L_{\odot}$) and density distribution, the abundance of gaseous water influences the two lines differently. The ground state line traces the amount of gaseous cold water, that is not frozen out onto grains in the outer regions due to cosmic rays and interstellar radiation heating. Previous observations (Boonman et al., 2002) estimate this abundance at $10^{-6} - 10^{-8}$ with respect to H_2 . The excited state traces the inner warm

region, but is still influenced by the cold water. Typical temperatures are $0.5-2$ K $km s^{-1}$ for the ground state and $0.1-3$ K $km s^{-1}$, which shows the large dependency of water on the abundance.

In a future publication (van Kempen et al., 2006, in prep) we present the results of a large parameter study on the emission of gas-phase rotational water lines, including the dependency of water on the luminosity of the protostar as well as the density distribution of the envelope. We conclude that an extensive knowledge about the local conditions of YSOs as well as a detailed modelling approach like we used, is essential in interpreting the future Herschel observations of water. ALMA will play an essential role in such observations if one wants to understand the role of water in the warm inner region. The resolution and wavelength coverage of ALMA will provide an excellent probe into these region with high-excited CO and water, providing strict constraints on the oxygen content and x-ray luminosity inside the high temperature regions of these protostellar systems.




Fig. 1. Physical ($T(r)$, $n(r)$) and chemical ($X(r)$) structure of a typical protostellar




Fig. 1: Ground state line of water for different abundances in warm region (X_0) and cold region (X_d) in integrated line fluxes (K)




Fig. 2: Excited line of water for different abundances. Note the dependence on both cold and warm gaseous water

Deuterated chemistry on amorphous water ice surfaces in the ISM

L. Kristensen, L. Amiaud, F. Dulieu, J-H. Fillion, A. Momeni, J. L. Lemaire

LERMA, UMR 8112 du CNRS, Observatoire de Paris et
Université de Cergy Pontoise, 5 mail Gay Lussac, 95031 Cergy Pontoise Cedex

We have shown (F. Dulieu et al talk and additional "Interaction of molecular hydrogen with Amorphous Water Ice Surfaces (ASW)" poster) that when both H_2 and D_2 are deposited at 10K on a 10ML amorphous water ice film, D_2 is occupying the more bound sites. When the temperature of the film increases, during a TPD (thermally programmed desorption) experiment, H_2 desorbs first leading then to a subsequent enrichment of the ice film in D_2 . Such a situation may occur in some cold dense regions of the ISM at the surface of water ice covered interstellar grains, during the long pre-collapse stage of the star formation process, then favouring a deuterated chemistry to take place. At a later stage, when the temperature increases again, the resulting deuterated molecules are then released in the gas phase and become observable. IRAS16293-2422 (ρ Ophiucus star forming region) is an example of such regions where deuterated species appear to be 10^3 to 10^4 times more abundant than the average galactic ratio.

Evaluation of spectral data for radioastronomy, Herschel and ALMA. Contribution to the improvement of this data.

Valerio Lattanzi¹, Brian Drouin², John Pearson², Adam Walters¹

¹ CESR, 9, avenue du Colonel Roche - Boite postale 4346 31028 Toulouse Cedex 4,
France

² Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena, California 91109, USA

Two spectroscopic databases are freely available through Internet and commonly used for radioastronomy, that of Cologne University (CDMS) and that of JPL. For CASSIS software (http://www.cesr.fr/~walters/web_cassis/) (which can be freely downloaded at <http://pc-126.cesr.fr/> for prediction and analysis of astrophysical spectra from radiotelescopes) both of these databases are combined for user-friendliness. In certain cases the same species may be found in both of these databases. We have hence carried out a rapid evaluation of these two databases (http://www.cesr.fr/~walters/web_cassis/Divers/Molec_tab.pdf) for astrophysical species and suggest a preference when necessary. This evaluation is gradually being improved and refined for priority species for which alternative data sources may be suggested. It will be used to provide templates with suggested database entries for CASSIS.

We have also started to contribute to the improvement in data available, especially in the new spectral region above 1 THz to be opened by Herschel and partially by terrestrial instruments in very good weather conditions. Measurements have recently been taken at JPL of the following species: HCO^+ , DCO^+ , H^{13}CO^+ , D^{13}CO^+ , CH_3D , CH_3CN , propane, HCCD and rovibrational transitions of DCCD.

Some examples of ongoing analysis will be given. In certain cases of species already measured at lower frequency, the data can be shifted by up to 100 MHz compared to earlier predictions. This shift is obviously very important for correctly identifying lines. Even small shifts of the order of a MHz can be important when accurate velocity information is to be obtained from the lineshape and Doppler shift.

Velocities of gas in star-forming regions: 70AU resolution and 1km/s results for OMC1.

H.D.Nissen¹, M.Gustafsson¹, J.L.Lemaire^{2,3}, Y.Clénet⁴, D.Rouan⁴, D.Field¹

¹Dept. of Physics and Astronomy, University of Aarhus, DK-8000 Aarhus C, Denmark

²Observatoire Paris-Meudon, LERMA and UMR8112 of the CNRS, Paris, France

³University of Cergy-Pontoise, Paris, France

⁴Observatoire Paris-Meudon, LESIA, Paris, France

The Orion Molecular Cloud (OMC1) ($D = 460$ pc) is the closest highly active massive star-forming region. It has accordingly been studied over a large range of wavelengths from the X-ray to the radio (e.g. [1-6]) and has become a type-site for studies of star formation. We report IR H_2 $v=1-0$ S(1) data for this region.

The data were obtained with the GriF instrument on the CHFT (see [7,8]). The novelty of these data is in the association of a high spatial resolution of $0.15''$ (70 AU) with velocity discrimination as good as 1 km s^{-1} (3 σ). Results show striking evidence for the presence of shocks within OMC1 through the clear association of gas motions of tens of km s^{-1} with bright IR emission of H_2 .

The study of 193 bright regions affords a detailed view of shocks within OMC1. Many new features are revealed, providing a new perspective on the nature of OMC1. Data are visualized in Fig. 3 in the poster.

The major conclusions of this study may be summarised as follows:

(i) 193 distinct bright H_2 emission features have been identified. 2/3 of these show clear radial velocity structure with well defined features of greater than 5 km s^{-1} , indicating that the emission is generated by shocks.

(ii) There are equal numbers of shocks in Peaks 1 and 2, implying a greater concentration of activity in Peak 2 which is less than half the size of Peak 1. Flows are found to be grouped more tightly in Peak 2. Blue-shifted flows are somewhat favoured over red-shifted, presumably due to dust obscuration. No velocities exceed $35\text{-}40 \text{ km s}^{-1}$, consistent with upper limits for C-type shocks in dense media.

(iii) In a distinct zone, lying south of Peak 1 and west of Peak 2, all flows are blue-shifted, with an average velocity of $-18 \pm 8 \text{ km s}^{-1}$, suggesting a collective outflow motion in this area.

(iv) Spatial correlation between maximum brightness of H_2 emission and maximum velocity reveals deviations typically of $\leq 0.5''$. These deviations are used to estimate the orientation angle of shocks, in the plane of the sky - but not their absolute direction. Orientations are found to be random in Peak 2 and with some preference for a NW direction in Peak 1, but show a strong preference for a NE-SW orientation in the outflow region (iii).

(v) In [8] it was noted that potential low mass star forming zones, identified through the presence of energetic flows in OMC1, showed higher densities and correspondingly higher energy flux, and are altogether less simply structured than more conventional Class 0 or 1 low mass isolated star formation regions. Here we identify a number of zones which show similar characteristics to those discussed in [8]. Data point in particular to the close physical association of numerous H_2 emission features. These associations may represent sites of multiple star formation characterised by episodic outbursts of activity.

(vi) The most striking feature is the blue-shifted outflow (iii). This is the IR counterpart to an outflow observed previously only in the radio [1,9]. The radio data show the presence of a flow, part of a general expansion, with a velocity of $18 \pm 6 \text{ km s}^{-1}$ in agreement with our value of $18 \pm 8 \text{ km s}^{-1}$. The origin of this outflow is uncertain; candidates are the massive radio sources I and n [10]. Both are situated near the origin of the blue-shifted outflow, and both have shown evidence of outflow along a NE-SW axis in radio observations [11,12].

(vii) There is a remarkable feature in the H_2 emission data with strong velocity components, which, on the basis of a VLT image, is strongly suggestive of a region in which a blue-shifted flow has burst from behind, from source I, through the material of OMC1 sending shock waves through the medium. The axis of the outflow passes right through this object.

References: [1] Genzel & Stutzki (1989) ARA&A, 27, 41. [2] Wright et al. (1996) ApJ, 469, 216. [3] O'Dell (2001) ARA&A, 39, 99. [4] Chrysostomou et al. (2000) MNRAS, 312, 103. [5] Feigelson et al. (2002) ApJ, 574, 258. [6] Doi et al. (2002) AJ, 124, 445. [7] Clénet et al. (2002) PASP, 114, 563. [8] Gustafsson et al. (2003) A&A, 411, 437. [9] Genzel et al. (1981) ApJ 244, 884. [10] Menten & Reid (1995) ApJ, 445, L157. [11] Greenhill et al. (2004) IAU Symp. 221. [12] Shuping et al. (2004) AJ, 128, 363.

Alcohol Chemistry In The Galactic Center. Hot Core Chemistry Without Hot Cores.

M. A. Requena-Torres¹, J. Martín-Pintado¹, A. Rodríguez-Franco¹,
S. Martín², N. J. Rodríguez-Fernández³ and P. de Vicente⁴

¹ Departamento de Astrofísica Molecular e Infrarroja, Instituto de Estructura de la Materia (CSIC), C/ Serrano 121, E--28006 Madrid, Spain

² Instituto de RadioAstronomía Milimétrica, Avda. Divina Pastora 7, Núcleo Central, E--18012 Granada, Spain

³ Observatoire de Bordeaux, L3AB(UMR 5804), CNRS/Université Bordeaux 1, BP 89, 2 rue de l'Observatoire, 33270 Floirac, France

⁴ Observatorio Astronómico Nacional, Centro Astronómico de Yebes, Apartado 148, 19080 Guadalajara, Spain

We have carried out a systematic study of CH₃OH, C₂H₅OH, (CH₃)₂O, HCOOCH₃, HCOOH, CH₃COOH, H₂CO and CS in different Galactic center (GC) clouds. The abundance of CH₃OH ranges from few 10⁻⁸ to 10⁻⁶ (see fig. 1). While the abundance ratio of the molecules relative to CH₃OH is basically independent of the CH₃OH abundance and only varies in a factor of ~4-8, the abundance ratio of CS to CH₃OH, not affected by grain chemistry, varies by a factor of 60. Our data are compared with observations of the same molecules in short-lived objects like the hot cores. The abundance and the abundance ratios of the complex molecules relative to CH₃OH in massive hot cores are similar to that found in the GC clouds. Alcohol chemistry is believed to be driven by gas phase reactions after evaporation of alcohols from grain mantles. Gas phase chemistry after ejection of alcohol from grains cannot explain the observed abundances and relative abundances of some molecules. Our data suggest that basically all the molecules related to alcohol chemistry could be produced on grain mantles and/or depleted from gas phase after their formation. This interpretation requires frequent shocks in the GC region to keep the high abundances of these molecules in gas phase and a rather uniform average composition of the icy grain mantles. The molecular clouds associated with the Sickle and the Thermal Radio Archers, which seem to be affected by UV radiation show lower abundances of C₂H₅OH relative to CH₃OH which could be explained by the combination shock ejection and photo dissociation.




Figure 1: Relative abundances of the observed molecules respect to that of CH₃OH as a function of the CH₃OH abundances.

Radiative lifetimes and association cross-sections of CO²⁺

T. Šedivcová, P. R. Ždánková and V. Špirko

Department of Molecular Modeling, Center for Biomolecules and Complex Molecular Systems Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic Flemingovo n 2, 166 10 Prague 6, Czech Republic

Despite the fact, all vibronic states of a dication are only metastable because of electrostatic repulsion, there are a couple of experimental measurements and also possibilities that these species could exist for a long time in the interstellar matter. As for CO²⁺ most of the lower electronic states have minima which are capable of supporting vibrational levels. Moreover, the levels that are trapped behind the barrier would classically be considered stable, but at the same time they lie in a continuum and so can lead to formation of the dication by radiative association.

This study is aimed to determinate the radiative lifetimes of vibrational levels in three lowest electronic states ³Π, ¹Σ⁺ and ¹Π and energy-dependent radiative association cross-section for the reaction C⁺(²Π_u) + O⁺(⁴Σ_u). The calculations are performed on the highly correlated icMRCI wavefunctions. These function have been used to calculate the potential energy (PE), spin-orbit coupling (SOC), dipole moment and transition moment functions for the lowest electronic states of CO²⁺ dication. Using PE and SOC functions, the positions and lifetimes of the corresponding vibronic states are evaluated by means of the complex coordinate method. Although, there are not many experimental results with which a comparison is possible, the calculated predissociating lifetimes are in qualitative accord with them.

Radiative lifetimes to v = 0 in terms of each one electronic state are shorter then predissociating lifetimes. So that the formation through radiative association is possible. As for radiative association cross-section, in spite of the transition dipole moment between singlet-triplet is spin-forbidden, singlet and triplet states in the {*ab initio*} calculations are not eigenstates of this CO²⁺ Hamiltonian. Transitions from ¹Σ⁺ and ¹Π to X³Π state occur because of mixing of states due to the spin-orbit of electronic states.

Reliability of the disk chemical modeling

A.I. Vasyunin¹, D. Semenov², A.M. Sobolev¹, Th. Henning²

¹ Ural State University, ul. Lenina 51, Yekaterinburg, 620083, Russia

² Max Planck Institute for Astronomy, Koenigstuhl 17, 69117 Heidelberg, Germany

We study how uncertainties of the rate coefficients of chemical reactions in the UMIST\95 database affect the modelled molecular abundances in a protoplanetary disk around a low-mass star. For that, we randomly varied the gas-phase reaction rates within their uncertainty limits and calculated sets of time-dependent abundances in several representative disk locations, using a gas-grain chemical model supplied with a set of surface reactions. All species are separated in different groups according to the sensitivity of their final abundances to the rate uncertainties. We found that such division depends on the disk location, since distinct processes control the chemical evolution of various disk regions. Typically, the modelled abundances of simple species are imprecise up to a half order of magnitude, and in general increase with the number of atoms in the molecule. We apply a straightforward sensitivity method introduced in [1] to analyze the correlations between the time-dependent abundances and varied rate values in order to identify those reactions whose rate uncertainties affect the resulting concentrations of the pre-selected, observationally important species to the largest extent. It is demonstrated that the rate coefficients of only a handful of chemical reactions need to be determined more accurately in order to improve significantly the reliability of disk astrochemical models, which might be a task for future laboratory studies and theoretical investigations.

References:


- 1) Vasyunin, A.I., Sobolev, A.M., Wiebe, D.S., & Semenov, D.A. (2004), *Astronomy Letters*, 30, 566
- 2) Wakelam, V., Selsis, F., Herbst, E., & Caselli, P. (2005), *Astronomy & Astrophysics*, 444, 883
- 3) Wakelam, V., Herbst, E., & Selsis, F. (2006), *ArXiv Astrophysics e-prints*, astro-ph/0601611

Rotational excitation of interstellar cyanoacetylene

Wernli M, Wiesenfeld L, **Faure A**, Valiron P

Laboratoire d'Astrophysique de Grenoble, UMR5571 CNRS, Université Joseph Fourier,
BP 53, 38041 Grenoble cedex 09, France

Cyanoacetylene (HC_3N), the simplest cyanopolyynes, is ubiquitous in space and is generally considered as an ideal probe of physical conditions in a variety of astronomical environments. We have computed rate coefficients for rotational excitation of HC_3N ($J=0-50$) by He and H_2 for kinetic temperatures lower than 100K. These rates have been obtained from extensive quantum and quasi-classical calculations using new high accuracy potential energy surfaces (PES). The symmetry of the PES is shown to induce strong collisional selection rules. As a first application, we present a simple steady-state population model that shows population inversions for the lowest HC_3N rotational sub-levels at H_2 densities in the range $\sim 10^3-10^5 \text{ cm}^{-3}$.



Rate coefficients for rotational excitation of HC_3N (initial $J_1=0$) by H_2 and He as a function of the final J_1 for a kinetic temperature of 20K. The quasi-classical results of Green & Chapman (1978) are compared to the present quantum calculations. The strong quantum interferences reflect the quasi-symmetry of the potential energy surfaces.

Planet Formation - Traced with ALMA

Sebastian Wolf¹, Dmitry Semenov¹, Gennaro D'Angelo²,
Amaya Moro-Martín³, and Yaroslav Pavlyuchenkov¹

¹ Max Planck Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany

² NASA Ames Research Center, Space Science and Astrobiology Division,
MS 245-3, Moffett Field, CA 94035, USA

³ Department of Astrophysical Sciences, Peyton Hall, Ivy Lane, Princeton University,
Princeton, NJ 08544, USA

Planets are expected to form in circumstellar disks, which are considered as the natural outcome of the protostellar evolution, at least in the case of low and medium mass stars. While a detailed picture of the evolution of the circumstellar environment, in particular of protoplanetary disks, has been developed, the planet formation process is still under discussion. Adequate constraints from observations are required in order to either verify or rule out existing hypotheses about proposed planet formation scenarios.

Numerical simulations convincingly demonstrate that high-resolution imaging performed with observational facilities that are already available or will become available in the near future will allow to trace the formation and early evolution of planets in circumstellar disks. Theoretical investigations show that the signatures of planet formation are usually much larger in size than the planet itself and thus more easily detectable. The specific result of the planet-disk interaction depends on the evolutionary stage of the disk. Primary signatures of planets embedded in disks are gaps in the case of young disks and characteristic asymmetric density patterns in more evolved debris disks.

Based on the example of the Butterfly star in the star-forming Taurus region, it will be discussed how observations with the Submillimeter Array (SMA) now or with the Atacama Large Millimeter Array (ALMA) in the near future can be used not only to constrain the global disk structure but also to characterize the process of the grain growth, which is indicative for the earliest stage of planet formation. Furthermore, simulations concerning the observation of disk structures tracing the formation of planets, such as large-scale vortices, spiral density waves, gaps, and inner holes, will be outlined. For the late stages of giant planet formation, we will discuss the numerical simulations which demonstrate that ALMA will even allow to trace young giant planets in nearby protoplanetary disks (see Fig. 1). Furthermore, we will show that molecular line observations of planet formation signatures with ALMA will allow us to study in detail the kinematics of gas and gas-to-dust ratio across the discussed features.




Figure 1: Simulation of ALMA observations of a disk with an embedded giant planet around a $0.5M_{\text{sun}}$ star (orbital radius: 5AU; assumed distance: 50 pc). The hot local environment of the $1M_{\text{jup}}$ planet is clearly seen as a bright region in the upper part of the low-density gap caused by the planet-disk-interaction. The disk mass amounts to $M_{\text{disk}} = 1.0 \times 10^{-2} M_{\text{sun}}$. The size of the combined beam is symbolized in the lower left corner. [based on Wolf & D'Angelo (2005) *ApJ*, 619, 1114]

List of Participants

Mike Alexandersen

Niels Bohr Institute
Copenhagen Ø
Denmark
mikea@fys.ku.dk

Cristina Apetrei

Physical Chemistry
University of Basel
Basel
Switzerland
cristina.apetrei@unibas.ch

Denise Bailly

LPPM
University Paris-Sud
Orsay
France
denise.bailly@ppm.u-psud.fr

Jeanette Bast

Astronomy
Stockholm University
Stockholm
Sweden
jeanette_bast@hotmail.com

Henrik Beuther

Max Planck Institute for Astronomy
Heidelberg
Germany
beuther@mpia.de

Ofer Biham

Racah Institute of Physics
The Hebrew University
Jerusalem
Israel
biham@phys.huji.ac.il

Rasmus Bjørk

Computational Astrophysics
Niels Bohr Institute
Copenhagen Ø
Denmark
rbjk@astro.ku.dk

Sandra Bruenken

CfA
Harvard University
Cambridge, MA
USA
sbruenken@cfa.harvard.edu

Franco Cataldo

Chemical Research Branch
Soc. Lupi Chemical Research Institute
Rome
Italy
cdcata@flashnet.it

Anja Andersen

Dark Cosmology Centre
Niels Bohr Institute
Copenhagen
Denmark
anja@astro.ku.dk

Rafael Bachiller

Observatorio Astronomico Nacional
Madrid
Spain
r.bachiller@oan.es

Jesper Bang

Department of Physics and Astronomy
University of Aarhus
Aarhus C
Denmark
jespergb@phys.au.dk

Lourdes Benito-Lopez

Univers Froid
Centre d'Etude Spatiale des Rayonnements
Toulouse
France
Lourdes.Benito@cesr.fr

Ludovic Biennier

Astrochimie Experimentale
Laboratoire PALMS UMR CNRS 6627
Rennes
France
ludovic.biennier@univ-rennes1.fr

Suzanne Bisschop

Raymond and Beverly Sackler Lab.
Leiden Observatory
RA Lieden
The Netherlands
bisschop@strw.leidenuniv.nl

Henrik Brink

Niels Bohr Institute
Copenhagen Ø
Denmark
henrikbrink@gmail.com

Piergiorgio Casavecchia

Department of Chemistry
University of Perugia
Perugia
Italy
piero@dyn.unipg.it

Cecilia Ceccarelli

Laboratoire d'Astrophysique de Grenoble
Grenoble
France
Cecilia.Ceccarelli@obs.ujf-grenoble.fr

Jose Cernicharo
DAMIR
Instituto de Estructura de la Materia
Madrid
Spain
cerni@damir.iem.csic.es

Sandip Chakrabarti
Theoretical Astrophysics
S.N. Bose National Centre for Basic
Sciences, Kolkata
India
chakraba@bose.res.in

Fabien Daniel
LERMA
Observatoire de Paris
Meudon
France
fabien.daniel@obspm.fr

Karine Demyk
PhLAM
Lille
France
karine.demyk@univ-lille1.fr

Zahia Djouadi
Institut d'Astrophysique Spatiale
Orsay
France
zahia.djouadi@ias.u-psud.fr

Francois Dulieu
LERMA
Université de Cergy Pontoise
Cergy-Pontoise
France
francois.dulieu@obspm.fr

Alexandre Faure
Laboratoire d'Astrophysique
Observatoire de Grenoble
Grenoble
France
afaure@obs.ujf-grenoble.fr

Carsten Frigaard
Department of Physics and Astronomy
University of Aarhus
Aarhus C
Denmark
carsten_frigaard@hotmail.com

Gary Fuller
School of Physics and Astronomy
University of Manchester
Manchester
United Kingdom
G.Fuller@manchester.ac.uk

Riccardo Cesaroni
Observatorio Astrofisico di Arcetri
INAF
Firenze
Italy
cesa@arcetri.astro.it

Claudia Comito
Max Planck Institute for Radioastronomie
Bonn
Germany
ccomito@mpifr-bonn.mpg.de

Frank De Lucia
Department of Physics
The Ohio State University
Columbus, OH
USA
fcd@mps.ohio-state.edu

Ewine van Dishoeck
Leiden University
Leiden
The Netherlands
ewine@strw.leidenuniv.nl

Marie-Lise Dubernet-Tuckey
LERMA
Observatory of Paris-Meudon
Meudon
France
marie-lise.dubernet@obspm.fr

David Edvardsson
Physical and Theoretical Chemistry Lab.
University of Oxford
Oxford
United Kingdom
david.edvardsson@chem.ox.ac.uk

David Field
Department of Physics & Astronomy
University of Aarhus
Aarhus C
Denmark
dfield@phys.au.dk

Asuncion Fuente
Observatorio Astronomico Nacional
Instituto Geografico Nacional
Madrid
Spain
a.fuente@oan.es

Wolf Dietrich Geppert
Department of Physics
Stockholm University
Stockholm
Sweden
wgeppert@hotmail.com

Thomas Giesen
Physikalisches Institut
University of Cologne
Cologne
Germany
giesen@ph1.uni-koeln.de

Antoine Gusdorf
Department of Physics
Durham University
Durham
United Kingdom
antoine.gusdorf@durham.ac.uk

Erik Herbst
Department of Physics
Ohio State University
Columbus Ohio
USA
herbst@mps.ohio-state.edu

Christine Joblin
CESR
CNRS
Toulouse
christine.joblin@cesr.fr

Antoine Jolly
LISA
University Paris 12
Paris
France
jolly@lisa.univ-paris12.fr

Tim van Kempen
Leiden Observatory
University of Leiden
Lieden
The Netherlands
kempen@strw.leidenuniv.nl

Valerio Lattanzi
CESR
Toulouse
France
valerio.lattanzi@cesr.fr

Francois Lique
LERMA
Observatoire de Meudon
Meudon
France
francois.lique@obspm.fr

Brenda Matthews
Herzberg Institute of Astrophysics
British Columbia
Canada
brenda.matthews@nrc-cnrc.gc.ca

Manuel Gonzalez
LUTH
Observatoire de Paris
Meudon
France
manuel.gonzalez@obspm.fr

Maiken Gustafsson
Department of Physics and Astronomy
University of Aarhus
Aarhus C
Denmark
maikeng@phys.au.dk

Liv Hornekaer
Department of Physics and Astronomy
University of Aarhus
Aarhus C
Denmark
liv@phys.au.dk

Jes Joergensen
Harvard-Smithsonian Centre for
Astrophysics
Cambridge, MA
USA
jjorgensen@cfa.harvard.edu

Eric Josselin
GRAAL
Universite Montpellier II
Montpellier
France
josselin@graal.univ-montp2.fr

Lars Egstrom Kristensen
Université de Cergy Pontoise
Cergy-Pontoise
France
egstrom@phys.au.dk

Jean-Louis Lemaire
LERMA
Observatoire de Paris
Meudon
France
jean-louis.lemaire@obspm.fr

Nigel Mason
Department of Physics & Astronomy
The Open University
Milton Keynes
United Kingdom
n.j.mason@open.ac.uk

Karl Menten
Millimeter and Submillimeter Astronomy
Max-Planck Institute for Radioastronomy
Bonn
Germany
kmenten@mpifr-bonn.mpg.de

Robert Moszynski

Department of Chemistry
University of Warsaw
Warsaw
Poland
robert.moszynski@tiger.chem.uw.edu.pl

Kaspar Nielsen

Computational Astrophysics
Niels Bohr Institute
Copenhagen
Denmark
kasparn@astro.ku.dk

Henrik Nissen

Department of Physics and Astronomy
University of Aarhus
Aarhus C
Denmark
hdn@phys.au.dk

Henrik Olofsson

Onsala Space Observatory
Onsala
Sweden
henrik@oso.chalmers.se

Berengere Parise

Max Planck Institute for Radioastronomy
Bonn
Germany
bparise@mpifr-bonn.mpg.de

Miguel Angel Requena-Torres

DAMIR
IEM-CSIC
Madrid
Spain
requena@damir.iem.csic.es

Nami Sakai

Department of Physics
University of Tokyo
Tokyo
Japan
nami@taurus.phys.s.u-tokyo.ac.jp

Stephan Schlemmer

Physikalisches Institut
University of Cologne
Cologne
Germany
schlemmer@ph1.uni-koeln.de

Tereza Sedivcova

Theoretical Chemistry
Institute of Organic Chemistry and
Biochemistry
Prague 6
Czech Republic
sedivcova@uochb.cas.cz

Holger Müller

Max Planck Institut für Radioastronomie
Bonn
Germany
hspm@ph1.uni-koeln.de

Jakob Holden Nielsen

Department of Physics and Astronomy
University of Aarhus
Aarhus C
Denmark
holden@phys.au.dk

Nagayoshi Ohashi

Institute of Astronomy and Astrophysics
Academia Sinica
Taipei
R. O. C.
ohashi@asiaa.sinica.edu.tw

Olja Panic

Leiden Observatory
University of Leiden
Leiden
The Netherlands
olja@strw.leidenuniv.nl

John Pearson

Jet Propulsion Laboratory
California
USA
John.C.Pearson@jpl.nasa.gov

Markus Roellig

KOSMA
University of Cologne
Cologne
Germany
roellig@ph1.uni-koeln.de

Peter Schilke

Max Planck Institute for Radioastronomie
Bonn
Germany
schilke@mpifr-bonn.mpg.de

Fredrik Schöier

Stockholm Observatory
Stockholm University
Stockholm
Sweden
fredrik@astro.su.se

Dmitry Semenov

Planet and Star Formation
Max Planck Institute for Astronomy
Heidelberg
Germany
semenov@mpia.de

Leonid Shirkov

Faculty of Chemistry, Quantum Chemistry
Warsaw University
Warsaw
Poland
leonid.shirkov@tiger.chem.uw.edu.pl

Sven Thorwirth

Max Planck Institut für Radioastronomie
Bonn
Germany
sthorwirth@mpifr-bonn.mpg.de

Michel Vervloet

LPPM
CNRS
Orsay
France
michel.vervloet@ppm.u-psud.fr

Nadine Wehres

Anion photoelectron spectroscopy lab
University of California
Berkeley
USA
wehres@strw.leidenuniv.nl

Laurent Wiesenfeld

Laboratoire d'Astrophysique
Observatoire de Grenoble
Grenoble
France
laurent.wiesenfeld@obs.ujf-grenoble.fr

Sebastian Wolf

Max Planck Institute for Astronomy
Heidelberg
Germany
swolf@mpia.de

Satoshi Yamamoto

Department of Physics
University of Tokyo
Tokyo
Japan
yamamoto@phys.s.u-tokyo.ac.jp

Bjarne Thomsen

Department of Physics and Astronomy
University of Aarhus
Aarhus C
Denmark
bt@phys.au.dk

Floris van der Tak

SRON National Institute for Space Research
Groningen
The Netherlands
vdtak@sron.rug.nl

Adam Walters

Univers Froid
Centre d'Etude Spatiale des Rayonnements
Toulouse
France
walters@cesr.fr

Michael Wernli

Laboratoire d'Astrophysique
Observatoire de Grenoble
Grenoble
France
michael.wernli@obs.ujf-grenoble.fr

Tom Wilson

ALMA
ESO
Garching
Germany
twilson@eso.org

Alywn Wootten

National Radio Astronomy Observatory
Charlottesville
USA
awootten@nrao.edu

John Yates Jr.

Department of Chemistry
University of Pittsburgh
Pittsburgh, PA
USA
jyates@pitt.edu

Index

- Acharyya, 52, 57, 58, 59
AlonsoAlbi, 53
Alsayed-Ali, 39
Amiaud, 63
Andersen, 29
Armengaud, 40
Bachiller, 20, 53
Bacmann, 56
Bailly, 54
Balucani, 42
Barzel, 55
Belloche, 12
Bergeat, 42
Bergin, 25
Bergman, 15
Bernath, 15
Beuther, 26
Biennier, 39
Biham, 55
Bisschop, 23, 56
Black, 15
Botinelli, 56
Brandenburg, 61
Brown, 33
Bruneleau, 40
Brünken, 33
Casavecchia, 42
Caselli, 60
Castets, 56
Cataldo, 44, 48
Caux, 56
Cazaux, 56
Ceccarelli, 21, 56, 60
Cernicharo, 28
Cesaroni, 27
Chakrabarti, S., 52, 57, 58, 59
Chakrabarti, S.K., 52, 57, 58, 59
Clénet, 65
Comito, 12, 56
Crapsi, 25
d'Hendecourt, 47
D'Angelo, 70
Das, 57, 58, 59
Davoisne, 47
de Lucia, 16
Demyk, 38
Djouadi, 47
Doty, 62
Drouin, 64
Dubernet, 17, 43
Dulieu, 46, 63
Ehlerding, 45
Faure, 69
Field, 61, 65
Fillion, 63
Frael, 40
Frisk, 15
Fuente, 53, 60
Fuller, 14
Geppert, 15, 45
Giesen, 31, 41
Godfrey, 33
Gustafsson, 61, 65
Hamberg, 45
Hasegawa, 15
Hellberg, 45
Helmich, 14, 56
Henning, 68
Herbst, 22
Hjalmarson, 15
Höfner, 29
Hogerheijde, 25, 62
Joblin, 40
Johnstone, 60
Jolly, 49
Jones, 47
Jørgensen, 23, 24, 25, 62
Kahane, 56
Källberg, 45
Kaminska, 45
Koning, 15
Kristensen, 63
Kwok, 15
Larsson, 15, 45
Lattanzi, 64
Lecacheux, 15
Lemaire, 61, 63, 65
Leonori, 42
Leroux, 47
Lewen, 31
Lipshtat, 55
Martín, 66
Martín-Pintado, 66
Matthews, 25
Menten, 12
Millar, 45
Momeni, 63
Moro-Martín, 70
Moszynski, 43
Müller, 12, 30
Natta, 53
Neri, 53, 60
Nissen, 65
Noguès, 40
Novotny, 39
Nummelin, 15
Ohashi, 19
Olberg, 15
Olofsson, 15
Österdahl, 45
Parise, 56
Pavlyuchenkov, 70

Pearson, 36, 64
Persson, 15
Petrucci, 42
Pijpers, 61
Pirali, 37, 54
Planesas, 53
Plume, 14
Requena-Torres, 66
Roberts, 45
Rodríguez-Fernández, 66
Rodríguez-Franco, 66
Rouan, 65
Rowe, 39
Sakai, N., 34
Sakai, T., 34
Sandqvist, 15
Schilke, 12, 13, 56
Schlemmer, 31
Schöier, 18
Šedivcová, 67
Segoloni, 42
Semaniak, 45
Semenov, 68, 70
Shirkov, 43
Simon, 40
Simonsson, 45
Snyder, 11
Sobolev, 68
Špirko, 67
Testi, 53
Thaddeus, 33
Thomas, 45
Thorwirth, 32
Tielens, 56
Travers, 39
Ugglas, 45
Valiron, 69
van Dishoeck, 10, 23, 56, 60, 62
van Kempen, 62
Vasyunin, 68
Vervloet, 37, 54
Vicente, 66
Volk, 15
Wakelam, 56
Walters, 17, 56, 64
Wiesenfeld, 69
Wilson, 11
Wirström, 15
Wolf, 70
Wootten, 35
Wyrowski, 60
Yamamoto, 34
Ždánská, 67
Zhaunerchyk, 45